

Indefinite affine hyperspheres admitting a pointwise symmetry

C. Scharlach*

February 2, 2008

Abstract

An affine hypersurface M is said to admit a pointwise symmetry, if there exists a subgroup G of $\text{Aut}(T_p M)$ for all $p \in M$, which preserves (pointwise) the affine metric h , the difference tensor K and the affine shape operator S . Here, we consider 3-dimensional indefinite affine hyperspheres, i. e. $S = \text{HId}$ (and thus S is trivially preserved). First we solve an algebraic problem. We determine the non-trivial stabilizers G of a traceless cubic form on a Lorentz-Minkowski space $\mathbb{R}_1^3$ under the action of the isometry group $SO(1, 2)$ and find a representative of each $SO(1, 2)/G$ -orbit. Since the affine cubic form is defined by h and K , this gives us the possible symmetry groups G and for each G a canonical form of K . Next, we classify hyperspheres admitting a pointwise G -symmetry for all non-trivial stabilizers G (apart from $\mathbb{Z}_2$). Besides well-known hyperspheres (for $\mathbb{Z}_2 \times \mathbb{Z}_2$ resp. $\mathbb{R}$ the hyperspheres have constant sectional curvature and Pick invariant $J < 0$ resp. $J = 0$) we obtain rich classes of new examples e.g. warped product structures of two-dimensional affine spheres (resp. quadrics) and curves. Moreover, we find a way to construct indefinite affine hyperspheres out of 2-dimensional quadrics or positive definite affine spheres.

Subject class: 53A15 (15A21, 53B30)

Keywords: 3-dimensional affine hyperspheres, indefinite affine metric, pointwise symmetry, $SO(1, 2)$ -action, stabilizers of a cubic form, affine differential geometry, affine spheres, reduction theorems, Calabi product of hyperbolic affine spheres

1 Introduction

Let M^n be a connected, oriented manifold. Consider an immersed hypersurface with relative normalization, i.e., an immersion $\varphi: M^n \rightarrow \mathbb{R}^{n+1}$

*Partially supported by the DFG-Project 'Geometric Problems and Special PDEs'

together with a transverse vector field ξ such that $D\xi$ has its image in φ_*T_xM . Equi-affine geometry studies the properties of such immersions under equi-affine transformations, i. e. volume-preserving linear transformations ($SL(n+1, \mathbb{R})$) and translations.

In the theory of nondegenerate equi-affine hypersurfaces there exists a canonical choice of transverse vector field ξ (unique up to sign), called the affine (Blaschke) normal, which induces a connection ∇ , a nondegenerate symmetric bilinear form h and a 1-1 tensor field S by

$$D_X Y = \nabla_X Y + h(X, Y)\xi, \quad (1)$$

$$D_X \xi = -SX, \quad (2)$$

for all $X, Y \in \mathcal{X}(M)$. The connection ∇ is called the induced affine connection, h is called the affine metric or Blaschke metric and S is called the affine shape operator. In general ∇ is not the Levi Civita connection $\hat{\nabla}$ of h . The difference tensor K is defined as

$$K(X, Y) = \nabla_X Y - \hat{\nabla}_X Y, \quad (3)$$

for all $X, Y \in \mathcal{X}(M)$. Moreover the form $h(K(X, Y), Z)$ is a symmetric cubic form with the property that for any fixed $X \in \mathcal{X}(M)$, $\text{trace } K_X$ vanishes. This last property is called the apolarity condition. The difference tensor K , together with the affine metric h and the affine shape operator S are the most fundamental algebraic invariants for a nondegenerate affine hypersurface (more details in Sec. 2). We say that M^n is indefinite, definite, etc. if the affine metric h is indefinite, definite, etc. For details of the basic theory of nondegenerate affine hypersurfaces we refer to [LSZ93] and [NS94].

Here we will restrict ourselves to the case of affine hyperspheres, i. e. the shape operator will be a (constant) multiple of the identity ($S = HId$). Geometrically this means that all affine normals pass through a fixed point or they are parallel. The abundance of affine hyperspheres dwarfs any attempts at a complete classification. Even with the restriction to locally strongly convex hyperspheres (i. e. h is positive definite) and low dimensions the class is simply too large to classify. In order to obtain detailed information one has therefore to revert to sub-classes such as the class of complete affine hyperspheres (see [LSZ93] and the references contained therein or i. e. [JL05] for a very recent result). Various authors have also imposed curvature conditions. In the case of constant curvature the classification is nearly finished (see [Vra00] and the references contained therein). In analogy to Chen's work, [Che93], a new curvature invariant for positive definite affine hyperspheres was introduced in [SSVV97]. A lower bound was given and, for $n = 3$, the classification of the extremal class was started. This classification was completed in [SV96], [KSV01] and [KV99]. The special (simple) form of the difference tensor K for this class is remarkable, actually it turns out that the hyperspheres admit a certain pointwise group symmetry [Vra04].

A hypersurface is said to admit a pointwise group symmetry if at every point the affine metric, the affine shape operator and the difference tensor are preserved under the group action. Necessarily the possible groups must be subgroups of the isometry group. The study of submanifolds which admit a pointwise group symmetry was initiated by Bryant in [Bry01] where he studied 3-dimensional Lagrangian submanifolds of $\mathbb{C}^3$, i. e. the isometry group is $\mathrm{SO}(3)$. Because of the similar basic invariants, Vrancken transferred the problem to 3-dimensional positive definite affine hyperspheres. A classification of 3-dimensional positive definite affine hyperspheres admitting pointwise symmetries was obtained in [Vra04] and then extended to positive definite hypersurfaces in [LS] (here the affine shape operator is non-trivial and thus no longer trivially preserved by isometries). Now, for the first time, we will consider the indefinite case, namely 3-dimensional indefinite affine hyperspheres.

We can assume that the affine metric has index two, i. e. the corresponding isometry group is the (special) Lorentz group $\mathrm{SO}(1, 2)$. Our question is the following: *What can we say about a three-dimensional indefinite affine hypersphere, for which there exists a non-trivial subgroup G of $\mathrm{SO}(1, 2)$ such that for every $p \in M$ and every $L \in G$:*

$$K(LX_p, LY_p) = L(K(X_p, Y_p)) \quad \forall X_p, Y_p \in T_p M.$$

In Section 2 we will state the basic formulas of (equi-)affine hypersurface-theory needed in the further classification. We won't need hypersurface-theory in Section 3 and 4, where we consider the group structure of $\mathrm{SO}(1, 2)$ and its action on cubic forms. In Section 3 we show that there exist six different normalforms of elements of $\mathrm{SO}(1, 2)$, depending on the eigenvalues and eigenspaces. We can always find an oriented basis of $\mathbb{R}_1^3$ such that every $L \in \mathrm{SO}(1, 2)$ has one (and only one) of the following matrix representations:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos t & -\sin t \\ 0 & \sin t & \cos t \end{pmatrix}, t \in (0, 2\pi) \setminus \{\pi\}, \quad \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad Id, \quad \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

all of these with respect to an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$, $\mathbf{t}$ timelike, $\mathbf{v}, \mathbf{w}$ spacelike, or

$$\begin{pmatrix} l & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \frac{1}{l} \end{pmatrix}, l \neq \pm 1, \quad \begin{pmatrix} 1 & -1 & -\frac{1}{2} \\ 0 & 1 & \frac{1}{2} \\ 0 & 0 & 1 \end{pmatrix},$$

with respect to a (LV)basis $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$, $\mathbf{e}, \mathbf{f}$ lightlike, $\mathbf{v}$ spacelike (Theorem 1).

Since we are interested in pointwise group symmetry, in Sec. 4 we study the nontrivial stabilizer of a traceless cubic form $\tilde{K}$ under the $\mathrm{SO}(1, 2)$ -action $\rho(L)(\tilde{K}) = \tilde{K} \circ L$. (cp. [Bry01] for the classification of the $\mathrm{SO}(3)$ -action). It turns out that the $\mathrm{SO}(1, 2)$ -stabilizer of a nontrivial traceless cubic form is isomorphic to either $\mathrm{SO}(2)$, $\mathrm{SO}(1, 1)$, $\mathbb{R}$, the group S_3 of order 6, $\mathbb{Z}_2 \times \mathbb{Z}_2$, $\mathbb{Z}_3$, $\mathbb{Z}_2$ or it is trivial (Theorem 2).

In the following we classify the indefinite affine hyperspheres which admit a pointwise $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry (Section 5), $\mathbb{R}$ -symmetry (Section 6),

SO(2)- or $\mathbb{Z}_3$ -symmetry (Section 7.1), S_3 -symmetry (Section 7.2) or SO(1, 1)-symmetry (Section 8). In case of $\mathbb{Z}_2 \times \mathbb{Z}_2$ resp. $\mathbb{R}$, we get the indefinite affine hyperspheres of constant sectional curvature with negative resp. vanishing Pick invariant J ; those with $J > 0$ are examples for $\mathbb{Z}_2$ -symmetry. The other classes are very rich, most of them are warped products of two-dimensional affine spheres ($\mathbb{Z}_3$) resp. quadrics (SO(2), SO(1, 1)), with a curve. Thus we get many new examples of 3-dimensional indefinite affine hyperspheres. Furthermore, we show how one can construct indefinite affine hyperspheres out of two-dimensional quadrics or positive definite affine spheres.

2 Basics of affine hypersphere theory

First we recall the definition of the affine normal ξ (cp. [NS94]). In equi-affine hypersurface theory on the ambient space $\mathbb{R}^{n+1}$ a fixed volume form $\det$ is given. A transverse vector field ξ induces a volume form θ on M by $\theta(X_1, \dots, X_n) = \det(\varphi_* X_1, \dots, \varphi_* X_n, \xi)$. Also the affine metric h defines a volume form ω_h on M , namely $\omega_h = |\det h|^{1/2}$. Now the affine normal ξ is uniquely determined (up to sign) by the conditions that $D\xi$ is everywhere tangential (which is equivalent to $\nabla\theta = 0$) and that

$$\theta = \omega_h. \quad (4)$$

Since we only consider 3-dimensional indefinite hyperspheres, i. e.

$$S = HId, \quad H = \text{const.} \quad (5)$$

we can fix the orientation of the affine normal ξ such that the affine metric has signature one. Then the sign of H in the definition of an affine hypersphere is an invariant.

Next we state some of the fundamental equations, which a nondegenerate hypersurface has to satisfy, see also [NS94] or [LSZ93]. These equations relate S and K with amongst others the curvature tensor R of the induced connection ∇ and the curvature tensor $\hat{R}$ of the Levi Civita connection $\hat{\nabla}$ of the affine metric h . There are the Gauss equation for ∇ , which states that:

$$R(X, Y)Z = h(Y, Z)SX - h(X, Z)SY,$$

and the Codazzi equation

$$(\nabla_X S)Y = (\nabla_Y S)X.$$

Also we have the total symmetry of the affine cubic form

$$C(X, Y, Z) = (\nabla_X h)(Y, Z) = -2h(K(X, Y), Z). \quad (6)$$

The fundamental existence and uniqueness theorem, see [Dil89] or [DNV90], states that given h , ∇ and S such that the difference tensor is symmetric

and traceless with respect to h , on a simply connected manifold M an affine immersion of M exists if and only if the above Gauss equation and Codazzi equation are satisfied.

From the Gauss equation and Codazzi equation above the Codazzi equation for K and the Gauss equation for $\widehat{\nabla}$ follow:

$$(\widehat{\nabla}_X K)(Y, Z) - (\widehat{\nabla}_Y K)(X, Z) = \frac{1}{2}(h(Y, Z)SX - h(X, Z)SY \\ - h(SY, Z)X + h(SX, Z)Y),$$

and

$$\widehat{R}(X, Y)Z = \frac{1}{2}(h(Y, Z)SX - h(X, Z)SY \\ + h(SY, Z)X - h(SX, Z)Y) - [K_X, K_Y]Z$$

If we define the Ricci tensor of the Levi-Civita connection $\widehat{\nabla}$ by:

$$\widehat{\text{Ric}}(X, Y) = \text{trace}\{Z \mapsto \widehat{R}(Z, X)Y\}. \quad (7)$$

and the Pick invariant by:

$$J = \frac{1}{n(n-1)}h(K, K), \quad (8)$$

then from the Gauss equation we immediately get for the scalar curvature $\widehat{\kappa} = \frac{1}{n(n-1)}(\sum_{i,j} h^{ij}\widehat{\text{Ric}}_{ij})$:

$$\widehat{\kappa} = H + J. \quad (9)$$

For an affine hypersphere the Gauss and Codazzi equations have the form:

$$R(X, Y)Z = H(h(Y, Z)X - h(X, Z)Y), \quad (10)$$

$$(\nabla_X H)Y = (\nabla_Y H)X, \quad \text{i. e.} \quad H = \text{const.}, \quad (11)$$

$$(\widehat{\nabla}_X K)(Y, Z) = (\widehat{\nabla}_Y K)(X, Z), \quad (12)$$

$$\widehat{R}(X, Y)Z = H(h(Y, Z)X - h(X, Z)Y) - [K_X, K_Y]Z. \quad (13)$$

Since H is constant, we can rescale φ such that $H \in \{-1, 0, 1\}$.

3 Normalforms in $\text{SO}(1, 2)$

We denote¹ by $\mathbb{R}_1^3$ the Pseudo-Euclidean vector space in which a non-degenerate indefinite bilinear form of index two is given. The bilinear form is called the inner product and denoted by $\langle \cdot, \cdot \rangle$. A basis $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ is called orthonormal (ONB) if

$$\langle \mathbf{t}, \mathbf{t} \rangle = -1, \quad \langle \mathbf{v}, \mathbf{v} \rangle = 1 = \langle \mathbf{w}, \mathbf{w} \rangle, \\ 0 = \langle \mathbf{t}, \mathbf{v} \rangle = \langle \mathbf{t}, \mathbf{w} \rangle = \langle \mathbf{v}, \mathbf{w} \rangle. \quad (14)$$

¹for the notation cp. [Gre67]

For a chosen ONB the inner product of two vectors is given by

$$\langle \mathbf{x}, \mathbf{y} \rangle = -x_t y_t + x_v y_v + x_w y_w, \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}_1^3. \quad (15)$$

A basis $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ is called a light-vector basis (LVB) if

$$\begin{aligned} \langle \mathbf{e}, \mathbf{e} \rangle &= 0 = \langle \mathbf{f}, \mathbf{f} \rangle, \langle \mathbf{e}, \mathbf{f} \rangle = 1, \\ \langle \mathbf{e}, \mathbf{v} \rangle &= 0 = \langle \mathbf{f}, \mathbf{v} \rangle, \langle \mathbf{v}, \mathbf{v} \rangle = 1. \end{aligned} \quad (16)$$

For a chosen LVB the inner product of two vectors is given by

$$\langle \mathbf{x}, \mathbf{y} \rangle = x_e y_f + x_f y_e + x_v y_v, \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}_1^3. \quad (17)$$

We want to consider the special Pseudo-Euclidean rotations $SO(1, 2)$, i. e. the linear transformations L of $\mathbb{R}_1^3$ which preserve the inner product and have determinant equal to one:

$$\langle L\mathbf{x}, L\mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle, \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}_1^3, \quad \det L = 1.$$

Depending on the eigenvalues and eigenspaces we get the following normal-forms of the elements of $SO(1, 2)$.

Theorem 1. *There exists a choice of an oriented basis of $\mathbb{R}_1^3$ such that every $L \in SO(1, 2)$ is of one (and only one) of the following types:*

1. (a) $A_t := \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos t & -\sin t \\ 0 & \sin t & \cos t \end{pmatrix}$, $t \in [0, 2\pi)$, $t \neq 0, \pi$,
for an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$, $\mathbf{t}$ timelike, $\mathbf{v}, \mathbf{w}$ spacelike,
eigenvalues: $\lambda_1 = 1$, eigenspaces $E(1) = \text{span}\{\mathbf{t}\}$ timelike.
- (b) $A_\pi := \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$, for an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$, as above,
eigenvalues: $\lambda_1 = 1$, $\lambda_{2,3} = -1$,
eigenspaces $E(1) = \text{span}\{\mathbf{t}\}$ timelike, $E(-1) = \text{span}\{\mathbf{v}, \mathbf{w}\}$, spacelike.
2. (a) $A_0 := \text{Id}$, for an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ as above or a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$,
eigenvalues: $\lambda_{1,2,3} = 1$, eigenspaces $E(1) = \mathbb{R}_1^3$.
- (b) $B := \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$, for an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$, as above, or a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$,
eigenvalues: $\lambda_{1,3} = -1$, $\lambda_2 = 1$,
eigenspaces $E(-1) = \text{span}\{\mathbf{t}, \mathbf{w}\} = \text{span}\{\mathbf{e}, \mathbf{f}\}$ timelike, $E(1) = \text{span}\{\mathbf{v}\}$, spacelike.
3. (a) $C_l := \begin{pmatrix} l & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \frac{1}{l} \end{pmatrix}$, $l \neq \pm 1$, for a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$,
eigenvalues: $\lambda_1 = l$, $\lambda_2 = 1$, $\lambda_3 = \frac{1}{l}$,
eigenspaces $E(l) = \text{span}\{\mathbf{e}\}$ lightlike, $E(1) = \text{span}\{\mathbf{v}\}$, spacelike,
 $E(\frac{1}{l}) = \text{span}\{\mathbf{f}\}$ lightlike.

(b) $C_1 := \begin{pmatrix} 1 & -1 & -\frac{1}{2} \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$, for a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$,
eigenvalues: $\lambda_{1,2,3} = 1$, eigenspaces $E(1) = \text{span}\{\mathbf{e}\}$ lightlike.

For the proof we will use the following

Lemma 1. Let $L \in \text{SO}(1, 2)$ with $L(\mathbf{e}) = l\mathbf{e}$, $l \neq 0$, for a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$. Since under L a LVB will be mapped to a LVB, the corresponding matrix must have the following form :

$$C_{l,m} = \begin{pmatrix} l & -lm & -l\frac{m^2}{2} \\ 0 & 1 & m \\ 0 & 0 & \frac{1}{l} \end{pmatrix}.$$

Proof. We will use the Notation: $\mathbf{e}' = L(\mathbf{e})$, $\mathbf{v}' = L(\mathbf{v})$, $\mathbf{f}' = L(\mathbf{f})$, and compute (16), using (17):

$$\begin{aligned} 0 &= \langle \mathbf{e}', \mathbf{v}' \rangle = lv'_f \Rightarrow v'_f = 0, \\ 1 &= \langle \mathbf{v}', \mathbf{v}' \rangle = (v'_v)^2 \Rightarrow v'_v = \varepsilon, \quad \varepsilon^2 = 1, \\ 1 &= \langle \mathbf{e}', \mathbf{f}' \rangle = lf'_f \Rightarrow f'_f = \frac{1}{l}, \\ 1 &= \det L = l\varepsilon \frac{1}{l} \Rightarrow \varepsilon = 1, \\ 0 &= \langle \mathbf{f}', \mathbf{v}' \rangle = v'_e \frac{1}{l} + f'_v \Rightarrow f'_v = -\frac{v'_e}{l}, \\ 0 &= \langle \mathbf{f}', \mathbf{f}' \rangle = 2f'_e \frac{1}{l} + \frac{(v'_e)^2}{2} \Rightarrow f'_e = -\frac{1}{l} \frac{(v'_e)^2}{2}. \end{aligned}$$

With $v'_e = -lm$ we obtain $C_{l,m}$. □

Proof of Thm. 1. Every $L \in \text{SO}(1, 2)$ must have a real eigenvalue since it is an automorphism of a three dimensional vector space. A corresponding eigenvector will be either timelike, spacelike or lightlike.

Let's consider first the case that we have (at least) a timelike eigenvector. We can choose an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$, such that $\mathbf{t}$ is the timelike eigenvector with eigenvalue $\varepsilon = \pm 1$. Then $\mathbf{v}, \mathbf{w}$ are spacelike and L restricted to $\mathbf{t}^\perp = \text{span}\{\mathbf{v}, \mathbf{w}\}$ is an isometry of $\mathbb{R}^2$, i. e. an Euclidean rotation.

If $\varepsilon = 1$, then L restricted to $\mathbf{t}^\perp$ is in $\text{SO}(2)$ (proper Euclidean rotation): In general we get no more real eigenvalues (case 1.(a)). If the restriction is a rotation by an angle of π , then we get the second eigenvalue -1 of multiplicity two. The eigenspace is $\mathbf{t}^\perp$ (and thus spacelike) (case 1.(b)). Finally, if the restriction (and thus L) is the identity map Id , every vector is an eigenvector and we also can choose a LVB (case 2.(a)).

If $\varepsilon = -1$, then L restricted to $\mathbf{t}^\perp$ is an improper Euclidean rotation of $\mathbb{R}^2$, thus it has eigenvalues 1 and -1 . We get the eigenvalue 1 with eigenspace $\text{span}\{\mathbf{v}\}$ (spacelike) and the eigenvalue -1 of multiplicity two

with eigenspace $\text{span}\{\mathbf{t}, \mathbf{w}\}$ (timelike). Since L restricted to $\text{span}\{\mathbf{t}, \mathbf{w}\}$ is equal to $-Id$, we also can choose a basis of two lightlike eigenvectors (case 2.(b)).

Next we will consider the case that we have (at least) a lightlike eigenvector. We can choose a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$, such that $\mathbf{e}$ is this lightlike eigenvector with eigenvalue $l \in \mathbb{R}$, $l \neq 0$. Since under L a LVB will be mapped to a LVB, the corresponding matrix must have the following form (cp. Lem. 1):

$$C_{l,m} = \begin{pmatrix} l & -lm & -l\frac{m^2}{2} \\ 0 & 1 & m \\ 0 & 0 & \frac{1}{l} \end{pmatrix}.$$

$C_{l,m}$ has the eigenvalues $\lambda_1 = l$, $\lambda_2 = 1$ and $\lambda_3 = \frac{1}{l}$. If $l \neq \pm 1$, we get three distinct real eigenvalues and the corresponding (1-dim.) eigenspaces are either lightlike ($\text{Eig}(l)$ and $\text{Eig}(\frac{1}{l})$) or spacelike ($\text{Eig}(1)$) (case 3.(a)). (Since we take an eigenvector basis, the LVB is up to the length of the lightlike eigenvectors uniquely determined.) If $l = -1$, $\lambda_1 = \lambda_3 = -1$, thus we have an eigenvalue of multiplicity two with eigenspace $\text{span}\{\mathbf{e}, \mathbf{f}\}$ (case 2.(b)). The case $l = 1$ is left, i. e. only one eigenvalue of multiplicity three: For $m = 0$ we obtain the identity map (case 2.(a)). For $m \neq 0$, the eigenspace $\text{Eig}(1) = \text{span}\{\mathbf{e}\}$ only is one-dimensional. To get a normalform of L we compute how $C_{1,m}$ changes if we choose another LVB $\{\mathbf{e}', \mathbf{v}', \mathbf{f}'\}$ (with the same orientation) where $\mathbf{e}'$ is an eigenvector of L ($\mathbf{e}' = a\mathbf{e}$). If we express $\{\mathbf{e}', \mathbf{v}', \mathbf{f}'\}$ in terms of $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ and compute (16), we obtain (cp. Lem. 1):

$$\begin{aligned} \mathbf{e}' &= a\mathbf{e}, \\ \mathbf{v}' &= v'_e \mathbf{e} + \mathbf{v}, \\ \mathbf{f}' &= -\frac{(v'_e)^2}{2a} \mathbf{e} - \frac{v'_e}{a} \mathbf{v} + \frac{1}{a} \mathbf{f}. \end{aligned}$$

The matrix representing L with respect to the new LVB, $C'_{1,m}$, has the form

$$C'_{1,m} = \begin{pmatrix} 1 & -\frac{m}{a} & -\frac{m^2}{2} \frac{1}{a^2} \\ 0 & 1 & \frac{m}{a} \\ 0 & 0 & 1 \end{pmatrix}. \text{ Thus } C'_{1,m} = C_{1,\frac{m}{a}}, \text{ and we can choose a LVB}$$

such that $\frac{m}{a} = 1$ (case 3.(b)). From the above computations we see that this LVB still isn't completely determined, we can choose v'_e arbitrary.

Finally the case is left that we have (at least) a spacelike eigenvector $\mathbf{v}$. The corresponding eigenvalue must be $\varepsilon = \pm 1$ and L restricted to $\mathbf{v}^\perp$ is an isometry of $\mathbb{R}_1^2$, i. e. a Pseudo-Euclidean rotation (boost). Thus it always has two real eigenvalues with one-dimensional eigenspaces. We can choose an eigenvector basis, which will be either an ONB or a LVB, and we get one of the following cases: 1.(b), 2.(a), 2.(b) or 3.(a) (cp. [Gre67], p. 273). $\square$

Remark. The choice of basis for the above normalforms is unique up to:

1. (a) the ONB is unique up to $\mathbf{t} \rightarrow \varepsilon \mathbf{t}, \mathbf{v}, \mathbf{w}$ up to a proper ($\varepsilon = 1$) or improper ($\varepsilon = -1$) Euclidean rotation in $\mathbb{R}^2$.
- (b) the ONB is unique up to $\mathbf{t} \rightarrow \varepsilon \mathbf{t}, \mathbf{v}, \mathbf{w}$ up to a proper ($\varepsilon = 1$) or improper ($\varepsilon = -1$) Euclidean rotation in $\mathbb{R}^2$.
2. (a) every ONB or LVB.
- (b) the ONB is unique up to $\mathbf{v} \rightarrow \varepsilon \mathbf{v}, \mathbf{t}, \mathbf{w}$ up to a proper ($\varepsilon = 1$) or improper ($\varepsilon = -1$) Pseudo-Euclidean rotation in $\mathbb{R}_1^2$,

$$\text{the LVB is unique up to } \begin{cases} \mathbf{v} \rightarrow \mathbf{v}, \\ \mathbf{e} \rightarrow a\mathbf{e}, \\ \mathbf{f} \rightarrow \frac{1}{a}\mathbf{f}, \end{cases} \text{ or } \begin{cases} \mathbf{v} \rightarrow -\mathbf{v}, \\ \mathbf{e} \rightarrow a\mathbf{f}, \\ \mathbf{f} \rightarrow \frac{1}{a}\mathbf{e}, \end{cases} \quad a \in \mathbb{R},$$

3. (a) the LVB is unique up to $\begin{cases} \mathbf{e} \rightarrow a\mathbf{e}, \\ \mathbf{f} \rightarrow \frac{1}{a}\mathbf{f}, \end{cases} \quad a \in \mathbb{R}.$ Under $\begin{cases} \mathbf{v} \rightarrow -\mathbf{v}, \\ \mathbf{e} \rightarrow a\mathbf{f}, \\ \mathbf{f} \rightarrow \frac{1}{a}\mathbf{e}, \end{cases}$,
 C_λ goes to $C_{\frac{1}{\lambda}}$.

$$(b) \text{ the LVB is unique up to } \begin{cases} \mathbf{e} \rightarrow \mathbf{e}, \\ \mathbf{v} \rightarrow b\mathbf{e} + \mathbf{v}, \\ \mathbf{f} \rightarrow -\frac{b^2}{2}\mathbf{e} - b\mathbf{v} + \mathbf{f}, \end{cases} \quad b \in \mathbb{R}.$$

Under $\begin{cases} \mathbf{e} \rightarrow a\mathbf{e}, \\ \mathbf{v} \rightarrow b\mathbf{e} + \mathbf{v}, \\ \mathbf{f} \rightarrow -\frac{b^2}{2a}\mathbf{e} - \frac{b}{a}\mathbf{v} + \frac{1}{a}\mathbf{f}, \end{cases}, C_1 \text{ goes to } C_{1, \frac{1}{a}}.$

4 Non-trivial $SO(1, 2)$ -stabilizers

Since we are interested in pointwise group symmetry, we study the nontrivial stabilizer of a traceless cubic form $\tilde{K}$ under the $SO(1, 2)$ -action $\rho(L)(\tilde{K}) = \tilde{K} \circ L$ resp.

$$\rho(L)(h(K(\cdot, \cdot), \cdot)) = h(K(L\cdot, L\cdot), L\cdot)$$

(cp. [Bry01] for the classification under the $SO(3)$ -action). We will use the following notation for the coefficients of the difference tensor K with respect to an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$:

$$K_{\mathbf{t}} = \begin{pmatrix} -a_1 & -a_2 & -a_3 \\ a_2 & a_4 & a_5 \\ a_3 & a_5 & a_1 - a_4 \end{pmatrix}, \quad K_{\mathbf{v}} = \begin{pmatrix} -a_2 & -a_4 & -a_5 \\ a_4 & a_6 & a_7 \\ a_5 & a_7 & a_2 - a_6 \end{pmatrix}, \quad (18)$$

$$K_{\mathbf{w}} = \begin{pmatrix} -a_3 & -a_5 & -(a_1 - a_4) \\ a_5 & a_7 & a_2 - a_6 \\ a_1 - a_4 & a_2 - a_6 & a_3 - a_7 \end{pmatrix},$$

resp. with respect to a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$:

$$K_{\mathbf{e}} = \begin{pmatrix} b_1 & b_4 & b_5 \\ b_2 & -2b_1 & b_4 \\ b_3 & b_2 & b_1 \end{pmatrix}, K_{\mathbf{v}} = \begin{pmatrix} b_4 & -2b_5 & b_6 \\ -2b_1 & -2b_4 & -2b_5 \\ b_2 & -2b_1 & b_4 \end{pmatrix}, K_{\mathbf{f}} = \begin{pmatrix} b_5 & b_6 & b_7 \\ b_4 & -2b_5 & b_6 \\ b_1 & b_4 & b_5 \end{pmatrix}. \quad (19)$$

We will prove the following theorem, stating not only the non-trivial stabilizers, but also give a normal form of K for each stabilizer.

Theorem 2. *Let $p \in M$ and assume that there exists a non-trivial element of $SO(1, 2)$ which preserves K . Then there exists an ONB resp. a LVB of $T_p M$ such that either*

1. $K = 0$, and this form is preserved by every isometry, or
2. $a_1 = 2a_4, a_4 > 0$, and all other coefficients vanish, this form is preserved by the subgroup $\{A_t, t \in \mathbb{R}\}$, isomorphic to $SO(2)$, or
3. $a_6 > 0$, and all other coefficients vanish, this form is preserved by the subgroup with generators $\langle A_{\frac{2\pi}{3}}, B \rangle$, isomorphic to S_3 , or
4. $a_1 = 2a_4, a_4 > 0, a_6 > 0$, and all other coefficients vanish, this form is preserved by the subgroup with generator $\langle A_{\frac{2\pi}{3}} \rangle$, isomorphic to $\mathbb{Z}_3$, or
5. $a_2, a_5 \in \mathbb{R}, a_6 \geq 0$, where $(a_2, a_6) \neq 0$, and all other coefficients vanish, this form is preserved by the subgroup with generator $\langle B \rangle$, isomorphic to $\mathbb{Z}_2$, or
6. $a_5 > 0$, and all other coefficients vanish, this form is preserved by the subgroup with generators $\langle A_\pi, B \rangle$, isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$, or
7. $a_1 > 0$ or $a_4 > 0, a_1 \neq 2a_4$, and all other coefficients vanish, this form is preserved by the subgroup with generator $\langle A_\pi \rangle$, isomorphic to $\mathbb{Z}_2$, or
8. $b_4 > 0$, and all other coefficients vanish, this form is preserved by the subgroup $\{C_l, l \in \mathbb{R} \setminus \{0\}\}$, isomorphic to $SO(1, 1)$, or
9. $b_7 > 0$, and all other coefficients vanish, this form is preserved by the subgroup $\left\{ \begin{pmatrix} 1 & -m & -\frac{m^2}{2} \\ 0 & 1 & m \\ 0 & 0 & 1 \end{pmatrix}, m \in \mathbb{R} \right\}$, isomorphic to $\mathbb{R}$.

To get ready for the proof, first we will find out what it means for K to be invariant under one element of $SO(1, 2)$. Some of the computations were done with the CAS Mathematica².

²http://www.math.tu-berlin.de/~schar/IndefSym_Stabilizers.html

Lemma 2. *Let $p \in M$ and assume, that K is invariant under the transformation $A_t \in \text{SO}(1, 2)$, $t \in (0, 2\pi)$. Then we get for the coefficients of K with respect to the corresponding ONB of $T_p M$:*

1. *if $t \neq \frac{2\pi}{3}, \pi, \frac{4\pi}{3}$, then $a_1 = 2a_4$, $a_4 \in \mathbb{R}$, and all other coefficients vanish,*
2. *if $t = \frac{2\pi}{3}$ or $t = \frac{4\pi}{3}$, then $a_1 = 2a_4$, $a_4, a_6, a_7 \in \mathbb{R}$, and all other coefficients vanish,*
3. *if $t = \pi$, then $a_1, a_4, a_5 \in \mathbb{R}$, and all other coefficients vanish,*

Proof. The proof is a straight forward computation, evaluating the equations $h(K(X, Y), Z) = h(K(A_t X, A_t Y), A_t Z)$ for $X, Y, Z \in \{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$. The computations were done with the CAS Mathematica. For all $t \in (0, 2\pi)$ we obtain from eq2 ($X, Y = \mathbf{t}$, $Z = \mathbf{v}$) and eq3 ($X, Y = \mathbf{t}$, $Z = \mathbf{w}$) that $a_2 = 0$ and $a_3 = 0$. If $t = \pi$, then eq7 ($X, Y, Z = \mathbf{v}$) and eq8 ($X, Y = \mathbf{v}$, $Z = \mathbf{w}$) give $a_6 = 0$ and $a_7 = 0$. If $t \neq \pi$, then eq4 ($X = \mathbf{t}$, $Y, Z = \mathbf{v}$) and eq5 ($X = \mathbf{t}$, $Y = \mathbf{v}$, $Z = \mathbf{w}$) lead to $a_5 = 0$ and $a_1 = 2a_4$. Now, for $t = \frac{2\pi}{3}$ or $t = \frac{4\pi}{3}$, all equations are true. Otherwise, only $a_6 = 0$ and $a_7 = 0$ solve eq7 and eq8. $\square$

Lemma 3. *Let $p \in M$ and assume, that K is invariant under the transformation $B \in \text{SO}(1, 2)$. Then we get for the coefficients of K with respect to the corresponding ONB of $T_p M$ that $a_2, a_5, a_6 \in \mathbb{R}$, and all other coefficients vanish.*

Proof. The computations were done with the CAS Mathematica, too. We obtain from eq1 ($X, Y, Z = \mathbf{t}$), eq3 ($X, Y = \mathbf{t}$, $Z = \mathbf{w}$), eq4 ($X = \mathbf{t}$, $Y, Z = \mathbf{v}$) and eq8 ($X, Y = \mathbf{v}$, $Z = \mathbf{w}$) that $a_1 = 0$, $a_3 = 0$, $a_4 = 0$ and $a_7 = 0$. $\square$

Lemma 4. *Let $p \in M$ and assume, that K is invariant under the transformation $C_l \in \text{SO}(1, 2)$, $l \in \mathbb{R} \setminus \{0, 1\}$. Then we get for the coefficients of K with respect to the corresponding LVB of $T_p M$:*

1. *if $l \neq -1$, then $b_4 \in \mathbb{R}$, and all other coefficients vanish,*
2. *if $l = -1$, then $b_2, b_4, b_6 \in \mathbb{R}$, and all other coefficients vanish.*

Proof. The computations were done with the CAS Mathematica, too. We obtain from eq1 ($X, Y, Z = \mathbf{e}$), eq3 ($X, Y = \mathbf{e}$, $Z = \mathbf{f}$), eq6 ($X = \mathbf{e}$, $Y, Z = \mathbf{f}$) and eq10 ($X, Y, Z = \mathbf{f}$) that $b_1 = 0$, $b_3 = 0$, $b_5 = 0$ and $b_7 = 0$. If $l \neq -1$, then eq2 ($X, Y = \mathbf{e}$, $Z = \mathbf{v}$) and eq9 ($X = \mathbf{v}$, $Y, Z = \mathbf{f}$) additionally give that $b_2 = 0$ and $b_6 = 0$. $\square$

Lemma 5. *Let $p \in M$ and assume, that K is invariant under the transformation $C_{1,m} \in \text{SO}(1,2)$, $m \in \mathbb{R} \setminus \{0\}$. Then we get for the coefficients of K with respect to the corresponding LVB of $T_p M$ that $b_7 \in \mathbb{R}$, and all other coefficients vanish.*

Proof. The computations were done with the CAS Mathematica, too. We obtain successively from eq2 ($X, Y = \mathbf{e}, Z = \mathbf{v}$), eq3 ($X, Y = \mathbf{e}, Z = \mathbf{f}$), eq5 ($X = \mathbf{e}, Y = \mathbf{v}, Z = \mathbf{f}$), eq6 ($X = \mathbf{e}, Y, Z = \mathbf{f}$), eq9 ($X = \mathbf{v}, Y, Z = \mathbf{f}$) and eq10 ($X, Y, Z = \mathbf{f}$) that $b_3 = 0$, $b_2 = 0$, $b_1 = 0$, $b_4 = 0$, $b_5 = 0$ and $b_6 = 0$. $\square$

In the following U denotes an arbitrary subgroup of $\text{SO}(1,2)$, which leaves K invariant. We want to find out to which extend K determines the properties of the elements of U .

Lemma 6. *If there exists $t \in (0, 2\pi)$, $t \neq \pi$, with $A_t \in U$, and $K \neq 0$, then we get for the timelike eigenvector $\mathbf{t}$ of A_t :*

1. for $t \neq \frac{2\pi}{3}$ and $t \neq \frac{4\pi}{3}$: $M\mathbf{t} = \mathbf{t}$ for all $M \in U$,
2. for $t = \frac{2\pi}{3}$ or $t = \frac{4\pi}{3}$: $M\mathbf{t} = \varepsilon\mathbf{t}$ for all $M \in U$,

Proof. Let $M \in U$. From Lem. 2 we know that

$$h(K(M\mathbf{t}, M\mathbf{t}), MY) = h(K(\mathbf{t}, \mathbf{t}), Y) = \begin{cases} -2a_4, & Y = \mathbf{t}, \\ 0, & Y = \mathbf{v} \text{ or } Y = \mathbf{w}. \end{cases}$$

Thus $K(M\mathbf{t}, M\mathbf{t}) = -2a_4 M\mathbf{t}$, furthermore $h(M\mathbf{t}, M\mathbf{t}) = h(\mathbf{t}, \mathbf{t}) = -1$. Now assume that $X = x\mathbf{t} + y\mathbf{v} + z\mathbf{w} \in T_p M$ has the same properties ($K(X, X) = -2a_4 X$ and $h(X, X) = -1$). This is equivalent to (cp. Lem. 2):

$$(-2x^2 - y^2 - z^2)a_4 = -2a_4 x, \quad (20)$$

$$2xya_4 + (y^2 - z^2)a_6 + 2yza_7 = -2a_4 y, \quad (21)$$

$$2xza_4 - 2yza_6 + (y^2 - z^2)a_7 = -2a_4 z, \quad (22)$$

$$-x^2 + y^2 + z^2 = -1. \quad (23)$$

If $a_4 \neq 0$, then (20) and (23) imply that $3x^2 - 2x - 1 = 0$ and $x^2 \geq 1$, this means $x = 1$ and $y = z = 0$. Thus $X = \mathbf{t}$ and $M\mathbf{t} = \mathbf{t}$.

If $a_4 = 0$, then (21) and (22) imply that

$$\begin{aligned} (y^2 - z^2)a_6 + 2yza_7 &= 0, \\ -2yza_6 + (y^2 - z^2)a_7 &= 0. \end{aligned}$$

The two equations, linear in a_6 and a_7 , only have a non-trivial solution if $y = z = 0$. With (23) we obtain that $X = \varepsilon\mathbf{t}$ and thus $M\mathbf{t} = \varepsilon\mathbf{t}$. $\square$

Lemma 7. *If there exists $l \in \mathbb{R}$, $l \neq 0, \pm 1$, with $C_l \in U$, and $K \neq 0$, then we get for the spacelike eigenvector $\mathbf{v}$ of C_l : $M\mathbf{v} = \mathbf{v}$ for all $M \in U$.*

Proof. Let $M \in U$. From Lem. 4 we know that

$$h(K(M\mathbf{v}, M\mathbf{v}), MY) = h(K(\mathbf{v}, \mathbf{v}), Y) = \begin{cases} -2a_4, & Y = \mathbf{v}, \\ 0, & Y = \mathbf{e} \text{ or } Y = \mathbf{f}. \end{cases}$$

Thus $K(M\mathbf{v}, M\mathbf{v}) = -2a_4M\mathbf{v}$, furthermore $h(M\mathbf{v}, M\mathbf{v}) = h(\mathbf{v}, \mathbf{v}) = 1$. Now assume that $X = x\mathbf{t} + y\mathbf{v} + z\mathbf{w} \in T_pM$ has the same properties ($K(X, X) = -2a_4X$ and $h(X, X) = 1$). This is equivalent to (cp. Lem. 4):

$$2xyb_4 = -2b_4x, \quad (24)$$

$$2(-y^2 + xz)b_4 = -2b_4y, \quad (25)$$

$$2yzb_4 = -2b_4z, \quad (26)$$

$$2xz + y^2 = 1. \quad (27)$$

Since $b_4 \neq 0$, (24) is equivalent to $x = 0$ or $y = -1$, and (26) is equivalent to $z = 0$ or $y = -1$. Now $y = -1$ in (25) gives $xy = 2$, which is a contradiction to (27). Thus $x = 0$ and $z = 0$. With (25) we obtain that $X = \mathbf{v}$ and thus $M\mathbf{v} = \mathbf{v}$. $\square$

Lemma 8. *If there exists $m \in \mathbb{R}$, $m \neq 0$, with $C_{1,m} \in U$, and $K \neq 0$, then we get for the lightlike eigenvector $\mathbf{e}$ of $C_{1,m}$: $M\mathbf{e} = \mathbf{e}$ for all $M \in U$.*

Proof. Let $M \in U$. From Lem. 5 we know that

$$K(X, X) = b_7z^2\mathbf{e} \text{ for all } X = x\mathbf{t} + y\mathbf{v} + z\mathbf{w} \in T_pM,$$

i. e. $\mathbf{e}$ is determined by K up to length: $\mathbf{e} = \frac{1}{h(K(\mathbf{f}, \mathbf{f}), \mathbf{f})}K(\mathbf{f}, \mathbf{f})$. From the invariance of K under M it follows that $M\mathbf{e} = \frac{1}{h(K(\mathbf{f}, \mathbf{f}), \mathbf{f})}K(M\mathbf{f}, M\mathbf{f}) = (h(M\mathbf{f}, \mathbf{e}))^2\mathbf{e}$. Since $1 = h(M\mathbf{f}, M\mathbf{e})$, we get now: $1 = h(M\mathbf{f}, h(M\mathbf{f}, \mathbf{e})^2\mathbf{e}) = h(M\mathbf{f}, \mathbf{e})^3$, thus $h(M\mathbf{f}, \mathbf{e}) = 1$. $\square$

Now we are ready for the proof of Thm. 2.

Proof of Theorem 2. For the proof we will consider several cases which are supposed to be exclusive. Let U be a maximal subgroup of $\text{SO}(1, 2)$ which leaves K invariant.

1. Case We assume that there exists $t \in (0, 2\pi)$, $t \neq \frac{2\pi}{3}, \pi, \frac{4\pi}{3}$, with $A_t \in U$. Thus there exists an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such that K has the form (Lem. 2): $a_1 = 2a_4$, $a_4 \in \mathbb{R}$ and all other coefficients vanish. If $a_4 = 0$, then $K = 0$ and $U = \text{SO}(1, 2)$. If $a_4 \neq 0$, then we know that $M\mathbf{t} = \mathbf{t}$ for all $M \in U$ (Lem. 6). We have seen in Sec. 3 that $M \in U$ must be of type $1(a)$, $1(b)$ or $2(a)$. Now let $N \in \text{SO}(1, 2)$ be of type $1(a)$, $1(b)$ or $2(a)$ with eigenvector $\mathbf{t}$. We can normalize simultaneously (i. e. find an ONB

such that both A_t and N have normalform) and we see that N leaves K invariant (Lem. 2). Thus $U = \{A_t, t \in \mathbb{R}\}$. Finally, if $a_4 < 0$, we can change the ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ and take instead $\{-\mathbf{t}, \mathbf{w}, \mathbf{v}\}$.

2. Case We assume that $A_{\frac{2\pi}{3}} \in U$ or $A_{\frac{4\pi}{3}} \in U$. Thus there exists an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such that K has the form (Lem. 2): $a_1 = 2a_4$, $a_4, a_6, a_7 \in \mathbb{R}$ and all other coefficients vanish. Without loss of generality a_6 and a_7 will not vanish both. Furthermore we know that $M\mathbf{t} = \varepsilon\mathbf{t}$ for all $M \in U$ (Lem. 6). We have seen in Sec. 3 that $M \in U$ must be of type $1(a)$, $1(b)$, $2(a)$ or $2(b)$. Now let $N \in \text{SO}(1, 2)$ be of type $1(a)$, $1(b)$, $2(a)$ or $2(b)$ with eigenvector $\mathbf{t}$. We can normalize simultaneously.

If $N = A_{\frac{2\pi}{3}}$, $N = A_{\frac{4\pi}{3}}$ or $N = \text{Id}$ (type $1(a)$ or $2(a)$), then it leaves K invariant (Lem. 2). If $N = A_\pi$ (type $1(b)$), then by Lem. 2 $a_6 = 0 = a_7$, which gives a contradiction. If $N = B$ (type $2(b)$), then by Lem. 3 $a_4 = 0 = a_7$ and a_6 is the only non-vanishing coefficient of K .

We get two possibilities for K and the corresponding maximal subgroup U . Either $a_6 \in \mathbb{R} \setminus \{0\}$ and all other coefficients of K vanish, and $U = \langle A_{\frac{2\pi}{3}}, B \rangle$, if necessary by a change of basis ($\{-\mathbf{t}, -\mathbf{v}, \mathbf{w}\}$) we can make sure that $a_6 > 0$. Or $a_1 = 2a_4$, $a_4, a_6, a_7 \in \mathbb{R}$ and all other coefficients vanish, and $U = \langle A_{\frac{2\pi}{3}} \rangle$. As before we can choose $\mathbf{t}$ such that $a_4 \geq 0$. A computation gives that under a change of ONB $\{\mathbf{t}^*, \mathbf{v}^*, \mathbf{w}^*\} = \{\mathbf{t}, \cos s\mathbf{v} + \sin s\mathbf{w}, -\sin s\mathbf{v} + \cos s\mathbf{w}\}$ we obtain for K (cp. (18)): $a_6^* = a_6 \cos(3s) + a_7 \sin(3s)$, $a_7^* = a_7 \cos(3s) - a_6 \sin(3s)$, i. e. there exists $s \in \mathbb{R}$ such that $a_7^* = 0$. We can change the sign of a_6 by switching from $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ to $\{\mathbf{t}, -\mathbf{v}, -\mathbf{w}\}$. Finally we see that $a_4 \neq 0$.

3. Case We assume that there exists $l \neq 0, \pm 1$, with $C_l \in U$. There exists a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ such that K has the form (Lem. 4): $b_4 \in \mathbb{R}$ and all other coefficients vanish. If $b_4 = 0$, then $K = 0$ and $U = \text{SO}(1, 2)$.

If $b_4 \neq 0$, then we know that $M\mathbf{v} = \mathbf{v}$ for all $M \in U$ (Lem. 7). We have seen in Thm. 1 that $M \in U$ must be of type $2(a)$, $2(b)$ or $3(a)$. Now let $N \in \text{SO}(1, 2)$ be of type $2(a)$, $2(b)$ or $3(a)$ with eigenvector $\mathbf{v}$. We can normalize simultaneously and we see that N leaves K invariant (Lem. 4, $B = C_{-1}$). Thus $U = \{C_l, l \in \mathbb{R} \setminus \{0\}\}$. Finally, if $b_4 < 0$, we can change the LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ to $\{\mathbf{f}, -\mathbf{v}, \mathbf{e}\}$ (cp. Rem. 3).

4. Case We assume that $C_1 \in U$. There exists a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ such that K has the form (Lem. 5): $b_7 \in \mathbb{R}$ and all other coefficients vanish. If $b_7 = 0$, then $K = 0$ and $U = \text{SO}(1, 2)$.

If $b_7 \neq 0$, then we know that $M\mathbf{e} = \mathbf{e}$ for all $M \in U$ (Lem. 8). We have seen in Thm. 1 that $M \in U$ must be of type $3(b)$. Now let $N \in \text{SO}(1, 2)$ be of type $3(b)$ with eigenvector $\mathbf{e}$, i. e. N has the form $C_{1,m}$, $m \in \mathbb{R}$. We see that N leaves K invariant (Lem. 5). Thus $U = \langle C_1 \rangle$ ($C_{1,m}C_{1,n} = C_{1,m+n}$). Finally, if $b_7 < 0$, we can change the LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ to $\{-\mathbf{e}, \mathbf{v}, -\mathbf{f}\}$ (cp. Rem. 3).

5. Case We assume that $A_\pi \in U$. There exists an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such

that K has the form (Lem. 2): $a_1, a_4, a_5 \in \mathbb{R}$ and all other coefficients vanish. Every $M \in U$ must be of type 1(b) or 2(b), otherwise we are in one of the foregoing cases.

a) Let $N \in \text{SO}(1, 2)$ be of type 1(b). There exists a spacelike eigenvector $\mathbf{w}$ such that $A_\pi \mathbf{w} = -\mathbf{w}$ and $N\mathbf{w} = -\mathbf{w}$, and we can choose an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such that A_π has normalform and $N = \begin{pmatrix} \cosh s & -\sinh s & 0 \\ \sinh s & -\cosh s & 0 \\ 0 & 0 & -1 \end{pmatrix}$. If we assume that K is invariant under N then we obtain for $s \neq 0$ that $K = 0$. The computations were done with the CAS Mathematica, too. We obtain from eq3 ($X, Y = \mathbf{t}, Z = \mathbf{w}$) and eq9 ($X = \mathbf{v}, Y, Z = \mathbf{w}$) that $a_5 = 0$ and $a_1 = a_4$, and from eq1 ($X, Y, Z = \mathbf{t}$) that $a_4 = 0$.

b) Let $N \in \text{SO}(1, 2)$ be of type 2(b). There exists a spacelike eigenvector $\mathbf{w}$ such that $A_\pi \mathbf{w} = -\mathbf{w}$ and $N\mathbf{w} = -\mathbf{w}$, and we can choose an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such that A_π has normalform and $N = \begin{pmatrix} -\cosh s & -\sinh s & 0 \\ \sinh s & \cosh s & 0 \\ 0 & 0 & -1 \end{pmatrix}$. If we assume that K is invariant under N then we obtain that $a_1 = 0 = a_4$. If $s \neq 0$, also $a_5 = 0$, i. e. $K = 0$. The computations were done with the CAS Mathematica, too. If $s \neq 0$, we get from eq3 ($X, Y = \mathbf{t}, Z = \mathbf{w}$) and eq9 ($X = \mathbf{v}, Y, Z = \mathbf{w}$) that $a_5 = 0$ and $a_1 = a_4$, and from eq1 ($X, Y, Z = \mathbf{t}$) that $a_4 = 0$. If $s = 0$, we get from eq1 ($X, Y, Z = \mathbf{t}$) and eq4 ($X = \mathbf{t}, Y, Z = \mathbf{v}$) that $a_1 = 0 = a_4$.

Summarized we got two different forms of K with corresponding maximal subgroups U : a) Either there exists an ONB such that $a_1, a_4, a_5 \in \mathbb{R}$, where $a_1 \neq 2a_4$ or $a_5 \neq 0$, and all other coefficients vanish, this form is preserved by $U = \langle A_\pi \rangle$. Since $K_{\mathbf{t}}$ is a symmetric operator on the positive definite space $\mathbf{t}^\perp$, we can diagonalize, then $a_5 = 0$. If necessary, we still can take $\{-\mathbf{t}, -\mathbf{v}, \mathbf{w}\}$ to get $a_1 > 0$ or $a_4 > 0$. b) In the other case there exists an ONB such that $a_5 \in \mathbb{R}$, $a_5 \neq 0$, and all other coefficients vanish, this form is preserved by $U = \langle A_\pi, B \rangle$. If $a_5 < 0$, we switch to the ONB $\{-\mathbf{t}, \mathbf{w}, \mathbf{v}\}$.

6. Case We assume that $B \in U$. There exists an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such that K has the form (Lem. 3): $a_2, a_5, a_6 \in \mathbb{R}$ and all other coefficients vanish. Every $M \in U$ must be of type 2(b), otherwise we are in one of the foregoing cases. Now let $N \in \text{SO}(1, 2)$ be of type 2(b). We can't normalize simultaneously. We only know that B and N both have two-dimensional timelike eigenspaces, which intersect in a line. This line g can be space-, time- or lightlike.

a) If g is spacelike, we can choose an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such that B has normalform and $N = \begin{pmatrix} -\cosh s & -\sinh s & 0 \\ \sinh s & \cosh s & 0 \\ 0 & 0 & -1 \end{pmatrix}$ (cp. case 5). If we assume that K is invariant under N then we obtain for $s \neq 0$ that $K = 0$. The computations were done with the CAS Mathematica, too. If $s \neq 0$, we get from eq3 ($X, Y = \mathbf{t}, Z = \mathbf{w}$) and eq9 ($X = \mathbf{v}, Y, Z = \mathbf{w}$) that $a_5 = 0$ and $a_2 = a_6$, and from eq1 ($X, Y, Z = \mathbf{t}$) that $a_6 = 0$.

b) If g is timelike, we can choose an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ such that B has

normalform and $N = \begin{pmatrix} -1 & 0 & 0 \\ 0 & \cos s & \sin s \\ 0 & \sin s & -\cos s \end{pmatrix}$. If we assume that K is invariant under N then we obtain for $s \neq 0, \frac{2\pi}{3}, \pi, \frac{4\pi}{3}$ that $K = 0$. For $s = \frac{2\pi}{3}, \pi, \frac{4\pi}{3}$ we are in one of the foregoing cases. The computations were done with the CAS Mathematica, too. If $s \neq 0, \pi$, we get from eq2 ($X, Y = \mathbf{t}, Z = \mathbf{v}$) and eq4 ($X = \mathbf{t}, Y, Z = \mathbf{v}$) that $a_2 = 0$ and $a_5 = 0$. If also $s \neq \frac{2\pi}{3}, \frac{4\pi}{3}$, then we get from eq10 ($X, Y, Z = \mathbf{w}$) that $a_6 = 0$.

c) If g is lightlike, we can choose a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ such that B has normalform and $N = \begin{pmatrix} -1 & -2m & 2m^2 \\ 0 & 1 & -2m \\ 0 & 0 & -1 \end{pmatrix}$ (use Lem. 1). If we assume that K is invariant under N then we obtain for $m \neq 0$ that $K = 0$. The computations were done with the CAS Mathematica, too. If $m \neq 0$, we get from eq5 ($X = \mathbf{t}, Y = \mathbf{v}, Z = \mathbf{w}$) and eq2 ($X, Y = \mathbf{t}, Z = \mathbf{v}$) that $b_6 = 0$ and $b_4 = 0$, and from eq1 ($X, Y, Z = \mathbf{t}$) that $b_2 = 0$.

Therefore we have that $U = \langle B \rangle$. If $a_6 < 0$, we can switch to $\{-\mathbf{t}, -\mathbf{v}, \mathbf{w}\}$. Since $K_{\mathbf{v}}$ is a symmetric operator on an indefinite space we can't always diagonalize. Thus we can't simplify K in general. $\square$

Remark. In the proof we only have used multilinear algebra. Thus the theorem stays true for an arbitrary $(1, 2)$ -tensor K on $\mathbb{R}_1^3$ with $\langle K(X, Y), Z \rangle$ totally symmetric and vanishing trace K_X .

5 Pointwise $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry

Let M^3 be a hypersphere admitting a pointwise $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry. According to Thm. 2, there exists for every $p \in M^3$ an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ of $T_p M^3$ such that

$$K(\mathbf{t}, \mathbf{t}) = 0, \quad K(\mathbf{t}, \mathbf{v}) = a_5 \mathbf{w}, \quad K(\mathbf{t}, \mathbf{w}) = a_5 \mathbf{v}, \quad (28)$$

$$K(\mathbf{v}, \mathbf{v}) = 0, \quad K(\mathbf{v}, \mathbf{w}) = -a_5 \mathbf{t}, \quad K(\mathbf{w}, \mathbf{w}) = 0. \quad (29)$$

Substituting this in Eq. (13), we obtain

$$\hat{R}(X, Y)Z = (H - a_5^2)(h(Y, Z)X - h(X, Z)Y). \quad (30)$$

Schur's Lemma implies that M^3 has constant sectional curvature, by the affine theorem egregium (9) we obtain that $\hat{\kappa} = H - a_5^2$ and $J = -a_5^2 < 0$. Affine hyperspheres with constant affine sectional curvature and nonzero Pick invariant were classified by Magid and Ryan [MR92]. They show in their main theorem that an affine hypersphere with Lorentz metric of constant curvature and nonzero Pick invariant is equivalent to an open subset of either $(x_1^2 + x_2^2)(x_3^2 + x_4^2) = 1$ or $(x_1^2 + x_2^2)(x_3^2 - x_4^2) = 1$. In both cases $\hat{\kappa} = 0$, i. e. $H = -J$. In the proof of the main theorem they explicitly show that only $(x_1^2 + x_2^2)(x_3^2 + x_4^2) = 1$ has negative Pick invariant and that K has normalform. This proves:

Theorem 3. *An affine hypersphere admits a pointwise $\mathbb{Z}_2 \times \mathbb{Z}_2$ -symmetry if and only if it is affine equivalent to an open subset of*

$$(x_1^2 + x_2^2)(x_3^2 + x_4^2) = 1.$$

For $(x_1^2 + x_2^2)(x_3^2 - x_4^2) = 1$ they compute that the only non-vanishing coefficient of K is a_2 . Thus it follows (Thm. 2):

Remark. The affine hypersphere $(x_1^2 + x_2^2)(x_3^2 - x_4^2) = 1$ admits a pointwise $\mathbb{Z}_2$ -symmetry.

6 Pointwise $\mathbb{R}$ -symmetry

Let M^3 be a hypersphere admitting a pointwise $\mathbb{R}$ -symmetry. According to Thm. 2, there exists for every $p \in M^3$ a LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ of $T_p M^3$ such that

$$K(\mathbf{e}, \mathbf{e}) = 0, \quad K(\mathbf{e}, \mathbf{v}) = 0, \quad K(\mathbf{e}, \mathbf{f}) = 0, \quad (31)$$

$$K(\mathbf{v}, \mathbf{v}) = 0, \quad K(\mathbf{v}, \mathbf{f}) = 0, \quad K(\mathbf{f}, \mathbf{f}) = b_7 \mathbf{e}. \quad (32)$$

Substituting this in Eq. (13), we obtain

$$\hat{R}(X, Y)Z = H(h(Y, Z)X - h(X, Z)Y). \quad (33)$$

Schur's Lemma implies that M^3 has constant sectional curvature, by the affine theorem egregium (9) we obtain that $\hat{\kappa} = H$ and $J = 0$. Affine hyperspheres with constant affine sectional curvature and zero Pick invariant were classified in [DMV00] (see Thm. 6.2 ($H = 0$), Thm. 7.2 ($H = 1$) and Thm. 8.2 ($H = -1$)). They are determined by a null curve in resp. $\mathbb{R}_1^3$, S_1^3 , H_1^3 , and a function along this curve (note that in the notion of [DMV00] (2) holds).

Theorem 4. *Let M^3 be an affine hypersphere admitting a pointwise $\mathbb{R}$ -symmetry. Then M^3 has constant sectional curvature $\hat{\kappa} = H$ and zero Pick invariant $J = 0$.*

Remark. A study of [DMV00] shows that an affine hypersphere admits a pointwise $\mathbb{R}$ -symmetry if and only if (2) holds (in their notations).

If (3) holds for an affine hypersphere with constant sectional curvature and zero Pick invariant, then it admits a pointwise $\mathbb{Z}_2$ -symmetry (cp. Thm. 2).

7 Pointwise $\text{SO}(2)$ -, S_3 - or $\mathbb{Z}_3$ -symmetry

Let M^3 be a hypersphere admitting a $\text{SO}(2)$ -, S_3 - or $\mathbb{Z}_3$ -symmetry. According to Thm. 2, there exists for every $p \in M^3$ an ONB $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ of $T_p M^3$

such that

$$\begin{aligned} K(\mathbf{t}, \mathbf{t}) &= -2a_4\mathbf{t}, & K(\mathbf{t}, \mathbf{v}) &= a_4\mathbf{v}, & K(\mathbf{t}, \mathbf{w}) &= a_4\mathbf{w}, \\ K(\mathbf{v}, \mathbf{v}) &= -a_4\mathbf{t} + a_6\mathbf{v}, & K(\mathbf{v}, \mathbf{w}) &= -a_6\mathbf{w}, & K(\mathbf{w}, \mathbf{w}) &= -a_4\mathbf{t} - a_6\mathbf{v}, \end{aligned}$$

where $a_4 > 0$ and $a_6 = 0$ in case of $SO(2)$ -symmetry, $a_4 = 0$ and $a_6 > 0$ for S_3 , and $a_4 > 0$ and $a_6 > 0$ for $\mathbb{Z}_3$.

We would like to extend the ONB locally. It is well known that $\widehat{\text{Ric}}$ (cp. (7)) is a symmetric operator and we compute (some of the computations in this section are done with the CAS Mathematica³):

Lemma 9. *Let $p \in M$ and $\{\mathbf{t}, \mathbf{v}, \mathbf{w}\}$ the basis constructed earlier. Then*

$$\begin{aligned} \widehat{\text{Ric}}(\mathbf{t}, \mathbf{t}) &= -2(H - 3a_4^2), & \widehat{\text{Ric}}(\mathbf{t}, \mathbf{v}) &= 0, \\ \widehat{\text{Ric}}(\mathbf{t}, \mathbf{w}) &= 0, & \widehat{\text{Ric}}(\mathbf{v}, \mathbf{v}) &= 2(H - a_4^2 + a_6^2), \\ \widehat{\text{Ric}}(\mathbf{v}, \mathbf{w}) &= 0, & \widehat{\text{Ric}}(\mathbf{w}, \mathbf{w}) &= 2(H - a_4^2 + a_6^2). \end{aligned}$$

Proof. The proof is a straight-forward computation using the Gauss equation (13). It follows e. g. that

$$\begin{aligned} \hat{R}(\mathbf{t}, \mathbf{v})\mathbf{t} &= H\mathbf{v} - K_{\mathbf{t}}(a_4\mathbf{v}) + K_{\mathbf{v}}(-2a_4\mathbf{t}) = H\mathbf{v} - a_4^2\mathbf{v} - 2a_4^2\mathbf{v} \\ &= (H - 3a_4^2)\mathbf{v}, \\ \hat{R}(\mathbf{t}, \mathbf{w})\mathbf{t} &= H\mathbf{w} - K_{\mathbf{t}}(a_4\mathbf{w}) + K_{\mathbf{w}}(-2a_4\mathbf{t}) = H\mathbf{w} - a_4^2\mathbf{w} - 2a_4^2\mathbf{w} \\ &= (H - 3a_4^2)\mathbf{w}, \\ \hat{R}(\mathbf{t}, \mathbf{v})\mathbf{w} &= -K_{\mathbf{t}}(-a_6\mathbf{w}) + K_{\mathbf{v}}(a_4\mathbf{w}) = 0. \end{aligned}$$

From this it immediately follows that

$$\widehat{\text{Ric}}(\mathbf{t}, \mathbf{t}) = -2(H - 3a_4^2)$$

and

$$\widehat{\text{Ric}}(\mathbf{t}, \mathbf{w}) = 0.$$

The other equations follow by similar computations. $\square$

We want to show that the basis, we have constructed at each point p , can be extended differentiably to a neighborhood of the point p such that, at every point, K with respect to the frame $\{T, V, W\}$ has the previously described form.

Lemma 10. *Let M^3 be an affine hypersphere in $\mathbb{R}^4$ which admits a pointwise $SO(2)$ -, S_3 - or $\mathbb{Z}_3$ -symmetry. Let $p \in M$. Then there exists an orthonormal*

³http://www.math.tu-berlin.de/~schar/IndefSym_typ234.html

frame $\{T, V, W\}$ defined in a neighborhood of the point p such that K is given by:

$$\begin{aligned} K(T, T) &= -2a_4T, & K(T, V) &= a_4V, & K(T, W) &= a_4W, \\ K(V, V) &= -a_4T + a_6V, & K(V, W) &= -a_6W, & K(W, W) &= -a_4T - a_6V, \end{aligned}$$

where $a_4 > 0$ and $a_6 = 0$ in case of $\text{SO}(2)$ -symmetry, $a_4 = 0$ and $a_6 > 0$ in case of S_3 -symmetry, and $a_4 > 0$ and $a_6 > 0$ in case of $\mathbb{Z}_3$ -symmetry.

Proof. First we want to show that at every point the vector $\mathbf{t}$ is uniquely defined (up to sign) and differentiable. We introduce a symmetric operator $\hat{A}$ by:

$$\widehat{\text{Ric}}(Y, Z) = h(\hat{A}Y, Z).$$

Clearly $\hat{A}$ is a differentiable operator on M . Since $2(H - 3a_4^2) \neq 2(H - a_4^2 + a_6^2)$, the operator has two distinct eigenvalues. A standard result then implies that the eigendistributions are differentiable. We take T a local unit vectorfield spanning the 1-dimensional eigendistribution, and local orthonormal vectorfields $\tilde{V}$ and $\tilde{W}$ spanning the second eigendistribution. If $a_6 = 0$, we can take $V = \tilde{V}$ and $W = \tilde{W}$.

As T is (up to sign) uniquely determined, for $a_6 \neq 0$ there exist differentiable functions a_4 , c_6 and c_7 , $c_6^2 + c_7^2 \neq 0$, such that

$$\begin{aligned} K(T, T) &= -2a_4T, & K(\tilde{V}, \tilde{V}) &= -a_4T + c_6\tilde{V} + c_7\tilde{W}, \\ K(T, \tilde{V}) &= a_4\tilde{V}, & K(\tilde{V}, \tilde{W}) &= c_7\tilde{V} - c_6\tilde{W}, \\ K(T, \tilde{W}) &= a_4\tilde{W}, & K(\tilde{W}, \tilde{W}) &= -a_4T - c_6\tilde{V} - c_7\tilde{W}. \end{aligned}$$

As we have shown in the proof of Thm. 2 (Case 2), we can always rotate $\tilde{V}$ and $\tilde{W}$ such that we obtain the desired frame. $\square$

Remark. It actually follows from the proof of the previous lemma that the vector field T is (up to sign) invariantly defined on M , and therefore the function a_4 , too. Since the Pick invariant (8) $J = \frac{1}{3}(-5a_4^2 + 2a_6^2)$, the function a_6 also is invariantly defined on the affine hypersphere M^3 .

In this section we always will work with the local frame constructed in the previous lemma. We denote the coefficients of the Levi-Civita connection with respect to this frame by:

$$\begin{aligned} \hat{\nabla}_T T &= a_{12}V + a_{13}W, & \hat{\nabla}_T V &= a_{12}T - b_{13}W, & \hat{\nabla}_T W &= a_{13}T + b_{13}V, \\ \hat{\nabla}_V T &= a_{22}V + a_{23}W, & \hat{\nabla}_V V &= a_{22}T - b_{23}W, & \hat{\nabla}_V W &= a_{23}T + b_{23}V, \\ \hat{\nabla}_W T &= a_{32}V + a_{33}W, & \hat{\nabla}_W V &= a_{32}T - b_{33}W, & \hat{\nabla}_W W &= a_{33}T + b_{33}V. \end{aligned}$$

We will evaluate first the Codazzi and then the Gauss equations ((12) and (13)) to obtain more informations.

Lemma 11. *Let M^3 be an affine hypersphere in $\mathbb{R}^4$ which admits a pointwise $\text{SO}(2)$ -, S_3 - or $\mathbb{Z}_3$ -symmetry and $\{T, V, W\}$ the corresponding ONB. If the symmetry group is*

SO(2), then $0 = a_{12} = a_{13} = a_{23} = a_{32}$, $a_{33} = a_{22}$ and
 $T(a_4) = -4a_{22}a_4$, $0 = V(a_4) = W(a_4)$,

S₃, then $0 = a_{12} = a_{13}$, $a_{23} = -3b_{13} = -a_{32}$, $a_{33} = a_{22}$ and
 $T(a_6) = -a_{22}a_6$, $V(a_6) = 3b_{33}a_6$, $W(a_6) = -3b_{23}a_6$,

$\mathbb{Z}_3$ and $\mathbf{a}_6^2 \neq 4\mathbf{a}_4^2$, then $0 = a_{12} = a_{13} = a_{23} = a_{32}$, $a_{33} = a_{22}$, $b_{13} = 0$,
 $T(a_4) = -4a_{22}a_4$, $0 = V(a_4) = W(a_4)$, and
 $T(a_6) = -a_{22}a_6$, $V(a_6) = 3b_{33}a_6$, $W(a_6) = -3b_{23}a_6$,

$\mathbb{Z}_3$ and $\mathbf{a}_6 = 2\mathbf{a}_4$, then $a_{12} = 2a_{22} = -2a_{33} = -b_{33}$,
 $a_{13} = -2a_{23} = -2a_{32} = b_{23}$, $b_{13} = 0$, and
 $T(a_4) = 0$, $V(a_4) = -4a_{22}a_4$, $W(a_4) = 4a_{23}a_4$,

Proof. An evaluation of the Codazzi equations (12) with the help of the CAS Mathematica leads to the following equations (they relate to eq1–eq6 and eq8–eq9 in the Mathematica notebook):

$$V(a_4) = -2a_{12}a_4, \quad T(a_4) = -4a_{22}a_4 + a_{12}a_6, \quad 0 = 4a_{23}a_4 + a_{13}a_6, \quad (34)$$

$$W(a_4) = -2a_{13}a_4, \quad 0 = 4a_{32}a_4 + a_{13}a_6, \quad T(a_4) = -4a_{33}a_4 - a_{12}a_6, \quad (35)$$

$$T(a_6) - V(a_4) = 3a_{12}a_4 - a_{22}a_6, \quad 0 = a_{13}a_4 + (a_{23} + 3b_{13})a_6, \quad (36)$$

$$W(a_4) = (a_{23} + a_{32})a_6, \quad W(a_6) = (-a_{23} + 3a_{32})a_4 - b_{23}a_6, \quad (37)$$

$$V(a_6) = (-a_{22} + a_{33})a_4 + 3b_{33}a_6,$$

$$T(a_6) = -a_{12}a_4 - a_{33}a_6, \quad W(a_4) = -3a_{13}a_4 + (-a_{32} + 3b_{13})a_6, \quad (38)$$

$$V(a_4) = (-a_{22} + a_{33})a_6, \quad W(a_6) = (3a_{23} - a_{32})a_4 - 3b_{23}a_6, \quad (39)$$

$$0 = (a_{23} - a_{32})a_4, \quad (40)$$

$$W(a_4) = -a_{13}a_4 + (a_{32} - 3b_{13})a_6. \quad (41)$$

From the first equation of (35) (we will use the notation (35).1) and (37).1 resp. (34).3 and (35).2 we get:

$$0 = 2a_{13}a_4 + (a_{23} + a_{32})a_6 \quad (42)$$

$$0 = 2(a_{23} + a_{32})a_4 + a_{13}a_6. \quad (43)$$

From (39).1) and (34).1 resp. (34).2 and (35).3 we get:

$$0 = -2a_{12}a_4 + 2(a_{22} - a_{33})a_6 \quad (44)$$

$$0 = 2(-a_{22} + a_{33})a_4 + a_{12}a_6. \quad (45)$$

We consider first the case, that $\mathbf{a}_6^2 \neq 4\mathbf{a}_4^2$. Then we obtain from the foregoing equations that $a_{13} = 0$, $a_{32} = -a_{23}$, $a_{12} = 0$ and $a_{33} = a_{22}$. Furthermore it

follows from (34).1 that $V(a_4) = 0$, from (34).2 that $T(a_4) = -4a_{22}a_4$ and from (34).3 that $a_{23}a_4 = 0$. Equation (35).1 becomes $W(a_4) = 0$, equation (36).2 $T(a_6) = -a_{22}a_6$ and (36).3 $(a_{23} + 3b_{13})a_6 = 0$. Finally equation (37).2 resp. 3 gives $W(a_6) = -3b_{23}a_6$ and $V(a_6) = 3b_{33}a_6$.

In case of $SO(2)$ -symmetry ($a_4 > 0$ and $a_6 = 0$) it follows that $a_{23} = 0$ and thus the statement of the theorem.

In case of S_3 -symmetry ($a_4 = 0$ and $a_6 > 0$) it follows that $a_{23} = -3b_{13}$ and thus the statement of the theorem.

In case of $\mathbb{Z}_3$ -symmetry ($a_4 > 0$ and $a_6 > 0$) it follows that $a_{23} = 0$ and $b_{13} = 0$ and thus the statement of the theorem.

In case that $a_6 = \pm 2a_4$ ($\neq 0$), we can choose V, W such that $\mathbf{a}_6 = 2\mathbf{a}_4$. Now equations (40), (34).3 and (36).3 lead to $a_{23} = a_{32}$, $a_{13} = -2a_{23}$ and $b_{13} = 0$. A combination of (34).2 and (35).3 gives $a_{12} = (a_{22} - a_{33})$, and then by equations (36).2, (34).1 and (34).2 that $a_{33} = -a_{22}$. Thus $T(a_4) = 0$ by (34).2, $V(a_4) = -4a_{22}a_4$ by (34).1 and $W(a_4) = 4a_{22}a_4$ by (35).1. Finally (37).2 and (35).1 resp. (37).3 and (34).1 imply that $b_{23} = -a_{23}$ resp. $b_{33} = -a_{22}$. $\square$

An evaluation of the Gauss equations (13) with the help of the CAS Mathematica leads to the following :

Lemma 12. *Let M^3 be an affine hypersphere in $\mathbb{R}^4$ which admits a pointwise $SO(2)$ -, S_3 - or $\mathbb{Z}_3$ -symmetry and $\{T, V, W\}$ the corresponding ONB. Then*

$$T(a_{22}) = -a_{22}^2 + a_{23}^2 + H - 3a_4^2, \quad (46)$$

$$T(a_{23}) = -2a_{22}a_{23}, \quad (47)$$

$$W(a_{22}) + V(a_{23}) = 0, \quad (48)$$

$$W(a_{23}) - V(a_{22}) = 0, \quad (49)$$

$$V(b_{13}) - T(b_{23}) = a_{22}b_{23} + (a_{23} + b_{13})b_{33}, \quad (50)$$

$$T(b_{33}) - W(b_{13}) = (a_{23} + b_{13})b_{23} - a_{22}b_{33}, \quad (51)$$

$$V(b_{33}) - W(b_{23}) = -a_{22}^2 - a_{23}^2 + 2a_{23}b_{13} + b_{23}^2 + b_{33}^2 + H + a_4^2 + 2a_6^2, \quad (52)$$

If the symmetry group is $\mathbb{Z}_3$, then $a_6^2 \neq 4a_4^2$.

Proof. The equations relate to eq11–eq13 and eq16 in the Mathematica notebook. If $a_6^2 = 4a_4^2$ ($\neq 0$), then we obtain by equations eq11.1 and eq12.3 resp. eq15.3 and eq12.3 that $2V(a_{22}) = -4a_{22}^2 - H + 3a_4^2$ resp. $2W(a_{23}) = 4a_{23}^2 + H - 3a_4^2$, thus $V(a_{22}) - W(a_{23}) = -2a_{22}^2 - 2a_{23}^2 - H + 3a_4^2$. This gives a contradiction to eq13.3, namely $V(a_{22}) - W(a_{23}) = -2a_{22}^2 - 2a_{23}^2 - H - 9a_4^2$. $\square$

7.1 Pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry

As the vector field T is globally defined, we can define the distributions $L_1 = \text{span}\{T\}$ and $L_2 = \text{span}\{V, W\}$. In the following we will investigate these distributions. For the terminology we refer to [Nöl96].

Lemma 13. *The distribution L_1 is autoparallel (totally geodesic) with respect to $\widehat{\nabla}$.*

Proof. From $\widehat{\nabla}_T T = a_{12}V + a_{13}W = 0$ (cp. Lemma 12) the claim follows immediately. $\square$

Lemma 14. *The distribution L_2 is spherical with mean curvature normal $U_2 = a_{22}T$.*

Proof. For $U_2 = a_{22}T \in L_1 = L_2^\perp$ we have $h(\widehat{\nabla}_{E_a} E_b, T) = h(E_a, E_b)h(U_2, T)$ for $E_a, E_b \in \{V, W\}$, and $h(\widehat{\nabla}_{E_a} U_2, T) = h(E_a(a_{22})T + a_{22}\widehat{\nabla}_{E_a} T, T) = 0$ (cp. Lemma 11 and (48), (49)). $\square$

Remark. a_{22} is independent of the choice of ONB $\{V, W\}$. It therefore is a globally defined function on M .

We introduce a coordinate function t by $\frac{\partial}{\partial t} := T$. Using the previous lemma, according to [PR93], we get:

Lemma 15. *(M^3, h) admits a warped product structure $M^3 = I \times_{e^f} N^2$ with $f : I \rightarrow \mathbb{R}$ satisfying*

$$\frac{\partial f}{\partial t} = a_{22}. \quad (53)$$

Proof. Prop. 3 in [PR93] gives the warped product structure with warping function $\lambda_2 : I \rightarrow \mathbb{R}$. If we introduce $f = \ln \lambda_2$, following the proof we see that $a_{22}T = U_2 = -\text{grad}(\ln \lambda_2) = -\text{grad } f$. $\square$

Lemma 16. *The curvature of N^2 is ${}^N K(N^2) = e^{2f}(H + 2a_6^2 + a_4^2 - a_{22}^2)$.*

Proof. From Prop. 2 in [PR93] we get the following relation between the curvature tensor $\hat{R}$ of the warped product M^3 and the curvature tensor $\tilde{R}$ of the usual product of pseudo-Riemannian manifolds ($X, Y, Z \in \mathcal{X}(M)$ resp. their appropriate projections):

$$\begin{aligned} \hat{R}(X, Y)Z &= \tilde{R}(X, Y)Z \\ &+ h(Y, Z)(\widehat{\nabla}_X U_2 - h(X, U_2)U_2) - h(\widehat{\nabla}_X U_2 - h(X, U_2)U_2, Z)Y \\ &- h(X, Z)(\widehat{\nabla}_Y U_2 - h(Y, U_2)U_2) + h(\widehat{\nabla}_Y U_2 - h(Y, U_2)U_2, Z)X \\ &+ h(U_2, U_2)(h(Y, Z)X - h(X, Z)Y) \end{aligned}$$

Now $\tilde{R}(X, Y)Z = {}^N \hat{R}(X, Y)Z$ for all $X, Y, Z \in TN^2$ and otherwise zero (cp. [O'N83], pg. 89, Corollary 58) and $K(N^2) = K(V, W) = \frac{h(-\hat{R}(V, W)V, W)}{h(V, V)h(W, W) - h(V, W)^2}$ (cp. [O'N83], pg. 77, the curvature tensor has the opposite sign). Since $h(X, Y) = e^{2f}h(X, Y)$ for $X, Y \in TN^2$, it follows that

$${}^N K(N^2) = e^{2f}h(-{}^N \hat{R}(V, W)V, W).$$

Finally we obtain by the Gauss equation (13) the last ingredient for the computation: $\hat{R}(V, W)V = -(H + 2a_6^2 + a_4^2)W$ (cp. the Mathematica notebook). $\square$

Summarized we have obtained the following structure equations (cp. (1), (2) and (3)), where $a_6 = 0$ in case of $\text{SO}(2)$ -symmetry resp. $b_{13} = 0$ in case of $\mathbb{Z}_3$ -symmetry:

$$D_T T = -2a_4 T - \xi, \quad (54)$$

$$D_T V = +a_4 V - b_{13} W, \quad (55)$$

$$D_T W = +b_{13} V + a_4 W, \quad (56)$$

$$D_V T = +(a_{22} + a_4) V, \quad (57)$$

$$D_W T = + (a_{22} + a_4) W, \quad (58)$$

$$D_V V = +a_6 V - b_{23} W + (a_{22} - a_4) T + \xi, \quad (59)$$

$$D_V W = +b_{23} V - a_6 W, \quad (60)$$

$$D_W V = - (b_{33} + a_6) W, \quad (61)$$

$$D_W W = +(b_{33} - a_6) V + (a_{22} - a_4) T + \xi, \quad (62)$$

$$D_X \xi = -HX, \quad (63)$$

The Codazzi and Gauss equations ((12) and (13)) have the form (cp. Lem. 11 and 12):

$$T(a_4) = -4a_{22}a_4, \quad 0 = V(a_4) = W(a_4) \quad (64)$$

$$T(a_6) = -a_{22}a_6, \quad V(a_6) = 3b_{33}a_6, \quad W(a_6) = -3b_{23}a_6, \quad (65)$$

$$T(a_{22}) = -a_{22}^2 + H - 3a_4^2, \quad V(a_{22}) = 0, \quad W(a_{22}) = 0, \quad (66)$$

$$V(b_{13}) - T(b_{23}) = a_{22}b_{23} + b_{13}b_{33}, \quad (67)$$

$$T(b_{33}) - W(b_{13}) = b_{13}b_{23} - a_{22}b_{33}, \quad (68)$$

$$V(b_{33}) - W(b_{23}) = -a_{22}^2 + b_{23}^2 + b_{33}^2 + H + a_4^2 + 2a_6^2, \quad (69)$$

where $a_6 = 0$ in case of $\text{SO}(2)$ -symmetry resp. $b_{13} = 0$ in case of $\mathbb{Z}_3$ -symmetry.

Our first goal is to find out how N^2 is immersed in $\mathbb{R}^4$, i. e. to find an immersion independent of t . A look at the structure equations (54) - (63) suggests to start with a linear combination of T and ξ .

We will solve the problem in two steps. First we look for a vector field X with $D_T X = \alpha X$ for some function α : We define $X := AT + \xi$ for some function A on M^3 . Then $D_T X = \alpha X$ iff $\alpha = -A$ and $\frac{\partial}{\partial t} A = -A^2 + 2a_4 A + H$, and $A := a_{22} - a_4$ solves the latter differential equation. Next we want to multiply X with some function β such that $D_T(\beta X) = 0$: We define a positive function β on $\mathbb{R}$ as the solution of the differential equation:

$$\frac{\partial}{\partial t} \beta = (a_{22} - a_4) \beta \quad (70)$$

with initial condition $\beta(t_0) > 0$. Then $D_T(\beta X) = 0$ and by (57), (63) and

(58) we get (since β , a_{22} and a_4 only depend on t):

$$D_T(\beta((a_{22} - a_4)T + \xi)) = 0, \quad (71)$$

$$D_V(\beta((a_{22} - a_4)T + \xi)) = \beta(a_{22}^2 - a_4^2 - H)V, \quad (72)$$

$$D_W(\beta((a_{22} - a_4)T + \xi)) = \beta(a_{22}^2 - a_4^2 - H)W. \quad (73)$$

To obtain an immersion we need that $\nu := a_{22}^2 - a_4^2 - H$ vanishes nowhere, but we only get:

Lemma 17. *The function $\nu = a_{22}^2 - a_4^2 - H$ is globally defined, $\frac{\partial}{\partial t}(e^{2f}\nu) = 0$ and ν vanishes identically or nowhere on $\mathbb{R}$.*

Proof. Since $0 = \frac{\partial}{\partial t} K(N^2) = \frac{\partial}{\partial t}(e^{2f}(2a_6^2 - \nu))$ (Lem. 16) and $\frac{\partial}{\partial t}(e^{2f}2a_6^2) = 0$ (cp. (65) and (53)), we get that $\frac{\partial}{\partial t}(e^{2f}\nu) = 0$. Thus $\frac{\partial}{\partial t}\nu = -2(\frac{\partial}{\partial t}f)\nu = -2a_{22}\nu$. $\square$

7.1.1 The first case: $\nu \neq 0$ on M^3

We may, by translating f , i.e. by replacing N^2 with a homothetic copy of itself, assume that $e^{2f}\nu = \varepsilon_1$, where $\varepsilon_1 = \pm 1$.

Lemma 18. *$\Phi := \beta((a_{22} - a_4)T + \xi): M^3 \rightarrow \mathbb{R}^4$ induces a proper affine sphere structure, say $\tilde{\phi}$, mapping N^2 into a 3-dimensional linear subspace of $\mathbb{R}^4$. $\tilde{\phi}$ is part of a quadric iff $a_6 = 0$.*

Proof. By (72) and (73) we have $\Phi_*(E_a) = \beta\nu E_a$ for $E_a \in \{V, W\}$. A further differentiation, using (59) (β and ν only depend on t), gives:

$$\begin{aligned} D_V\Phi_*(V) &= \beta\nu D_VV \\ &= \beta\nu((a_{22} - a_4)T + a_6V - b_{23}W + \xi) \\ &= a_6\Phi_*(V) - b_{23}\Phi_*(W) + \nu\Phi \\ &= a_6\Phi_*(V) - b_{23}\Phi_*(W) + \varepsilon_1e^{-2f}\Phi. \end{aligned}$$

Similarly, we obtain the other derivatives, using (60) - (62), thus:

$$D_V\Phi_*(V) = a_6\Phi_*(V) - b_{23}\Phi_*(W) + e^{-2f}\varepsilon_1\Phi, \quad (74)$$

$$D_V\Phi_*(W) = b_{23}\Phi_*(V) - a_6\Phi_*(W), \quad (75)$$

$$D_W\Phi_*(V) = -(b_{33} + a_6)\Phi_*(W), \quad (76)$$

$$D_W\Phi_*(W) = (b_{33} - a_6)\Phi_*(V) + e^{-2f}\varepsilon_1\Phi, \quad (77)$$

$$D_{E_a}\Phi = \beta e^{-2f}\varepsilon_1 E_a. \quad (78)$$

The foliation at $f = f_0$ gives an immersion of N^2 to M^3 , say π_{f_0} . Therefore, we can define an immersion of N^2 to $\mathbb{R}^4$ by $\tilde{\phi} := \Phi \circ \pi_{f_0}$, whose structure equations are exactly the equations above when $f = f_0$. Hence, we know

that $\tilde{\phi}$ maps N^2 into $\text{span}\{\Phi_*(V), \Phi_*(W), \Phi\}$, an affine hyperplane of $\mathbb{R}^4$ and $\frac{\partial}{\partial t}\Phi = 0$ implies $\Phi(t, v, w) = \tilde{\phi}(v, w)$.

We can read off the coefficients of the difference tensor $K^{\tilde{\phi}}$ of $\tilde{\phi}$ (cf. (1) and (3)): $K^{\tilde{\phi}}(\tilde{V}, \tilde{V}) = a_6\tilde{V}$, $K^{\tilde{\phi}}(\tilde{V}, \tilde{W}) = -a_6\tilde{W}$, $K^{\tilde{\phi}}(\tilde{W}, \tilde{W}) = -a_6\tilde{V}$, and see that $\text{trace}(K^{\tilde{\phi}})_X$ vanishes. The affine metric introduced by this immersion corresponds with the metric on N^2 . Thus $\varepsilon_1\tilde{\phi}$ is the affine normal of $\tilde{\phi}$ and $\tilde{\phi}$ is a proper affine sphere with mean curvature ε_1 . Finally the vanishing of the difference tensor characterizes quadrics. $\square$

Our next goal is to find another linear combination of T and ξ , this time only depending on t . (Then we can express T in terms of ϕ and some function of t .)

Lemma 19. *Define $\delta := HT + (a_{22} + a_4)\xi$. Then there exist a constant vector $C \in \mathbb{R}^4$ and a function $a(t)$ such that*

$$\delta(t) = a(t)C.$$

Proof. Using (57) resp. (58) and (63) we obtain that $D_V\delta = 0 = D_W\delta$. Hence δ depends only on the variable t . Moreover, we get by (54), (66), (64) and (63) that

$$\begin{aligned} \frac{\partial}{\partial t}\delta &= D_T(HT + (a_{22} + a_4)\xi) \\ &= H(-2a_4T - \xi) + (-a_{22}^2 + H - 3a_4^2 - 4a_{22}a_4)\xi - (a_{22} + a_4)HT \\ &= -(3a_4 + a_{22})(HT + (a_{22} + a_4)\xi) \\ &= -(3a_4 + a_{22})\delta. \end{aligned}$$

This implies that there exists a constant vector C in $\mathbb{R}^4$ and a function $a(t)$ such that $\delta(t) = a(t)C$. $\square$

Notice that for an improper affine hypersphere ($H = 0$) ξ is constant and parallel to C . Combining $\tilde{\phi}$ and δ we obtain for T (cp. Lem. 18 and 19) that

$$T(t, v, w) = -\frac{a}{\nu}C + \frac{1}{\beta\nu}(a_{22} + a_4)\tilde{\phi}(v, w). \quad (79)$$

In the following we will use for the partial derivatives the abbreviation $\varphi_x := \frac{\partial}{\partial x}\varphi$, $x = t, v, w$.

Lemma 20.

$$\begin{aligned} \varphi_t &= -\frac{a}{\nu}C + \frac{\partial}{\partial t}\left(\frac{1}{\beta\nu}\right)\tilde{\phi}, \\ \varphi_v &= \frac{1}{\beta\nu}\tilde{\phi}_v, \\ \varphi_w &= \frac{1}{\beta\nu}\tilde{\phi}_w. \end{aligned}$$

Proof. As by (70) and Lem. 17 $\frac{\partial}{\partial t} \frac{1}{\beta\nu} = \frac{1}{\beta\nu}(a_{22} + a_4)$, we obtain the equation for $\varphi_t = T$ by (79). The other equations follow from (72) and (73). $\square$

It follows by the uniqueness theorem of first order differential equations and applying a translation that we can write

$$\varphi(t, v, w) = \tilde{a}(t)C + \frac{1}{\beta\nu}(t)\tilde{\phi}(v, w)$$

for a suitable function $\tilde{a}$ depending only on the variable t . Since C is transversal to the image of $\tilde{\phi}$ (cp. Lem. 18 and 19, $\nu \neq 0$), we obtain that after applying an equiaffine transformation we can write: $\varphi(t, v, w) = (\gamma_1(t), \gamma_2(t)\phi(v, w))$, in which $\tilde{\phi}(v, w) = (0, \phi(v, w))$. Thus we have proven the following:

Theorem 5. *Let M^3 be an indefinite affine hypersphere of $\mathbb{R}^4$ which admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry. Let $a_{22}^2 - a_4^2 \neq H$ for some $p \in M^3$. Then M^3 is affine equivalent to*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (t, v, w) \mapsto (\gamma_1(t), \gamma_2(t)\phi(v, w)),$$

where $\phi : N^2 \rightarrow \mathbb{R}^3$ is a (positive definite) elliptic or hyperbolic affine sphere and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve.

Moreover, if M^3 admits a pointwise $SO(2)$ -symmetry then N^2 is either an ellipsoid or a two-sheeted hyperboloid.

We want to investigate the conditions imposed on the curve γ . For this we compute the derivatives of φ :

$$\begin{aligned} \varphi_t &= (\gamma'_1, \gamma'_2\phi), & \varphi_v &= (0, \gamma_2\phi_v), & \varphi_w &= (0, \gamma_2\phi_w), \\ \varphi_{tt} &= (\gamma''_1, \gamma''_2\phi), & \varphi_{tv} &= (0, \gamma'_2\phi_v), & \varphi_{tw} &= (0, \gamma'_2\phi_w), \\ \varphi_{vv} &= (0, \gamma_2\phi_{vv}), & \varphi_{vw} &= (0, \gamma_2\phi_{vw}), & \varphi_{ww} &= (0, \gamma_2\phi_{ww}). \end{aligned} \quad (80)$$

Furthermore we have to distinguish if M^3 is proper ($H = \pm 1$) or improper ($H = 0$).

First we consider the case that M^3 is proper, i. e. $\xi = -H\varphi$. An easy computation shows that the condition that ξ is a transversal vector field, namely $0 \neq \det(\varphi_t, \varphi_v, \varphi_w, \xi) = -\gamma_2^2(\gamma_1\gamma'_2 - \gamma'_1\gamma_2)\det(\phi_v, \phi_w, \phi)$, is equivalent to $\gamma_2 \neq 0$ and $\gamma_1\gamma'_2 - \gamma'_1\gamma_2 \neq 0$. To check the condition that ξ is the Blaschke normal (cp. (4)), we need to compute the Blaschke metric h , using (1), (80), (74)–(77) and the notation $r, s \in \{v, w\}$ and g for the Blaschke metric of ϕ :

$$\begin{aligned} \varphi_{tt} &= \dots \varphi_t + \frac{\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2}{H(\gamma_1\gamma'_2 - \gamma'_1\gamma_2)}\xi, \\ \varphi_{tr} &= \text{tang.} \\ \varphi_{rs} &= \text{tang.} - \frac{\gamma'_1\gamma_2}{H(\gamma_1\gamma'_2 - \gamma'_1\gamma_2)}\varepsilon_1 g\left(\frac{\partial}{\partial r}, \frac{\partial}{\partial s}\right)\xi. \end{aligned}$$

We obtain that $\det h = h_{tt}(h_{vv}h_{ww} - h_{vw}^2) = \frac{\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2}{H^3(\gamma_1\gamma'_2 - \gamma'_1\gamma_2)^3}(\gamma'_1)^2\gamma_2^2 \det g$.

Thus (4) is equivalent to $\gamma_2^4(\gamma_1\gamma'_2 - \gamma'_1\gamma_2)^2 \det(\phi_v, \phi_w, \phi)^2 = |\frac{\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2}{(\gamma_1\gamma'_2 - \gamma'_1\gamma_2)^3}(\gamma'_1)^2\gamma_2^2 \det g|$.

Since ϕ is a definite proper affine sphere with normal $-\varepsilon_1\phi$, we can again use (4) to obtain

$$\xi = -H\varphi \iff \gamma_2^2|\gamma_1\gamma'_2 - \gamma'_1\gamma_2|^5 = |\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2|(\gamma'_1)^2 \neq 0.$$

From the computations above (g is positive definite) also it follows that φ is indefinite iff either $H \operatorname{sign}(\gamma_1\gamma'_2 - \gamma'_1\gamma_2) = \operatorname{sign}(\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2) = \operatorname{sign}(\gamma'_1\gamma_2\varepsilon_1)$ or $-H \operatorname{sign}(\gamma_1\gamma'_2 - \gamma'_1\gamma_2) = \operatorname{sign}(\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2) = \operatorname{sign}(\gamma'_1\gamma_2\varepsilon_1)$.

Next we consider the case that M^3 is improper, i. e. ξ is constant. By Lem. 19 ξ is parallel to C and thus transversal to ϕ . Hence we can apply an affine transformation to obtain $\xi = (1, 0, 0, 0)$. An easy computation shows that the condition that ξ is a transversal vector field, namely $0 \neq \det(\varphi_t, \varphi_v, \varphi_w, \xi) = -\gamma_2^2\gamma'_2 \det(\phi_v, \phi_w, \phi)$, is equivalent to $\gamma_2 \neq 0$ and $\gamma'_2 \neq 0$. To check the condition that ξ is the Blaschke normal (cp. (4)) we need to compute the Blaschke metric h , using (1), (80), (74)–(77) and the notation $r, s \in \{v, w\}$ and g for the Blaschke metric of ϕ :

$$\begin{aligned} \varphi_{tt} &= \dots \varphi_t - \frac{\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2}{\gamma'_2}\xi, \\ \varphi_{tr} &= \text{tang.} \\ \varphi_{rs} &= \text{tang.} + \frac{\gamma'_1\gamma_2}{\gamma'_2}\varepsilon_1 g\left(\frac{\partial}{\partial r}, \frac{\partial}{\partial s}\right)\xi. \end{aligned}$$

We obtain that $\det h = h_{tt}(h_{vv}h_{ww} - h_{vw}^2) = -\frac{\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2}{(\gamma'_2)^3}(\gamma'_1)^2\gamma_2^2 \det g$. Thus (4) is equivalent to $\gamma_2^4(\gamma'_2)^2 \det(\phi_v, \phi_w, \phi)^2 = |\frac{\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2}{(\gamma'_2)^3}(\gamma'_1)^2\gamma_2^2 \det g|$. Since ϕ is a definite proper affine sphere with normal $-\varepsilon_1\phi$, we can again use (4) to obtain

$$\xi = (1, 0, 0, 0) \iff \gamma_2^2|\gamma'_2|^5 = |\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2|(\gamma'_1)^2 \neq 0.$$

From the computations above also it follows that φ is indefinite iff either $-\operatorname{sign}(\gamma'_2) = \operatorname{sign}(\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2) = \operatorname{sign}(\gamma'_1\gamma_2\varepsilon_1)$ or $\operatorname{sign}(\gamma'_2) = \operatorname{sign}(\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2) = \operatorname{sign}(\gamma'_1\gamma_2\varepsilon_1)$.

Now we are ready for the converse.

Theorem 6. *Let $\phi : N^2 \rightarrow \mathbb{R}^3$ be a positive definite elliptic or hyperbolic affine sphere (with mean curvature $\varepsilon_1 = \pm 1$), and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve such that $\varphi(t, v, w) = (\gamma_1(t), \gamma_2(t)\phi(v, w))$ defines a 3-dimensional indefinite affine hypersphere. Then $\varphi(N^2 \times I)$ admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry.*

(i) *If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_2^2|\gamma_1\gamma'_2 - \gamma'_1\gamma_2|^5 = \operatorname{sign}(\gamma'_1\gamma_2\varepsilon_1)(\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2)(\gamma'_1)^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.*

(ii) If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_2^2 |\gamma_2'|^5 = \text{sign}(\gamma_1' \gamma_2 \varepsilon_1) (\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2') (\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite improper affine hypersphere.

Proof. We already have shown that φ defines a 3-dimensional indefinite proper resp. improper affine hypersphere. To prove the symmetry we need to compute K . By assumption, ϕ is an affine sphere with Blaschke normal $\xi^\phi = -\varepsilon_1 \phi$. For the structure equations (1) we use the notation $\phi_{rs} = \phi \Gamma_{rs}^u \phi_u - g_{rs} \varepsilon_1 \phi$, $r, s, u \in \{v, w\}$. Furthermore we introduce the notation $\alpha = \gamma_1 \gamma_2' - \gamma_1' \gamma_2$. Note that $\alpha' = \gamma_1 \gamma_2'' - \gamma_1'' \gamma_2'$.

(i) Using (80), we get the structure equations (1) for φ :

$$\begin{aligned}\varphi_{tt} &= \frac{\alpha'}{\alpha} \varphi_t + \frac{\gamma_1' \gamma_2'' - \gamma_1'' \gamma_2'}{H \alpha} \xi, \\ \varphi_{tr} &= \frac{\gamma_2'}{\gamma_2} \varphi_r, \\ \varphi_{rs} &= \phi \Gamma_{rs}^u \phi_u - g_{rs} \varepsilon_1 \frac{\gamma_1 \gamma_2'}{\alpha} \varphi_t - g_{rs} \varepsilon_1 \frac{\gamma_1' \gamma_2}{H \alpha} \xi.\end{aligned}$$

We compute K using (6) and obtain:

$$\begin{aligned}(\nabla_{\varphi_t} h)(\varphi_r, \varphi_s) &= \left(\left(\frac{\gamma_1 \gamma_2'}{\alpha} \right)' \frac{\alpha}{\gamma_1 \gamma_2} - 2 \frac{\gamma_2'}{\gamma_2} \right) h(\varphi_r, \varphi_s), \\ (\nabla_{\varphi_r} h)(\varphi_t, \varphi_t) &= 0,\end{aligned}$$

implying that K_{φ_t} restricted to the space spanned by φ_v and φ_w is a multiple of the identity. Taking T in direction of φ_t , we see that φ_v and φ_w are orthogonal to T . Thus we can construct an ONB $\{T, V, W\}$ with V, W spanning $\text{span}\{\varphi_v, \varphi_w\}$ such that $a_1 = 2a_4$, $a_2 = a_3 = a_5 = 0$. By the considerations in Sec. 4 we see that φ admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry.

(ii) The proof runs completely analog. $\square$

7.1.2 The second case: $\nu \equiv 0$ and $H \neq 0$ on M^3

Next, we consider the case that $H = a_{22}^2 - a_4^2$ and $H \neq 0$ on M^3 . It follows that $a_{22} \neq \pm a_4$ on M^3 .

We already have seen that M^3 admits a warped product structure. The map Φ we have constructed in Lemma 18 will not define an immersion (cp. (72) and (73)). Anyhow, for a fixed point t_0 , we get from (59) - (62), (72) and (73), using the notation $\tilde{\xi} = (a_{22} - a_4)T + \xi$:

$$\begin{aligned}D_V V &= a_6 V - b_{23} W + \tilde{\xi}, \\ D_V W &= b_{23} V - a_6 W, \\ D_W V &= -(b_{33} + a_6) W, \\ D_W W &= (b_{33} - a_6) V + \tilde{\xi}, \\ D_{E_a} \tilde{\xi} &= 0, \quad E_a \in \{V, W\}.\end{aligned}$$

Thus, if v and w are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(t_0, v, w)$ as a positive definite improper affine sphere in a 3-dimensional linear subspace.

Moreover, we see that this improper affine sphere is a paraboloid provided that $a_6(t_0, v, w)$ vanishes identically. From the differential equations (65) determining a_6 , we see that this is the case exactly when a_6 vanishes identically, i.e. when M^3 admits a pointwise $SO(2)$ -symmetry.

After applying a translation and a change of coordinates, we may assume that

$$\varphi(t_0, v, w) = (v, w, f(v, w), 0),$$

with affine normal $\tilde{\xi}(t_0, v, w) = (0, 0, 1, 0)$. To obtain T at t_0 , we consider (57) and (58) and get that

$$D_{E_a}(T - (a_{22} + a_4)\varphi) = 0, \quad E_a, E_b \in \{V, W\}.$$

Evaluating at $t = t_0$, this means that there exists a constant vector C , transversal to $\text{span}\{V, W, \xi\}$, such that $T(t_0, v, w) = (a_{22} + a_4)(t_0)\varphi(t_0, v, w) + C$. Since $a_{22} + a_4 \neq 0$ everywhere, we can write:

$$T(t_0, v, w) = \alpha_1(v, w, f(v, w), \alpha_2), \quad (81)$$

where $\alpha_1, \alpha_2 \neq 0$ and we applied an equiaffine transformation so that $C = (0, 0, 0, \alpha_1\alpha_2)$. To obtain information about $D_T T$ we have that $D_T T = -2a_4 T - \xi$ (cp. (54)) and $\xi = \tilde{\xi} - (a_{22} - a_4)T$ by the definition of $\tilde{\xi}$. Also we know that $\tilde{\xi}(t_0, v, w) = (0, 0, 1, 0)$ and by (71) - (73) that $D_X(\beta\tilde{\xi}) = 0$, $X \in \mathcal{X}(M)$. Taking suitable initial conditions for the function β ($\beta(t_0) = 1$), we get that $\beta\tilde{\xi} = (0, 0, 1, 0)$ and finally the following vector valued differential equation:

$$D_T T = (a_{22} - 3a_4)T - \frac{1}{\beta}(0, 0, 1, 0).$$

Solving this differential equation, taking into account the initial conditions (81) at $t = t_0$, we get that there exist functions δ_1 and δ_2 depending only on t such that

$$T(t, u, v) = (\delta_1(t)v, \delta_1(t)w, \delta_1(t)(f(v, w) + \delta_2(t)), \alpha_2\delta_1(t)),$$

where $\delta_1(t_0) = \alpha_1$, $\delta_2(t_0) = 0$, $\delta_1'(t) = (a_{22} - 3a_4)\delta_1(t)$ and $\delta_2'(t) = \delta_1^{-1}(t)\beta^{-1}(t)$. As $T(t, v, w) = \frac{\partial \varphi}{\partial t}(t, v, w)$ and $\varphi(t_0, v, w) = (v, w, f(v, w), 0)$ it follows by integration that

$$\varphi(t, v, w) = (\gamma_1(t)v, \gamma_1(t)w, \gamma_1(t)f(v, w) + \gamma_2(t), \alpha_2(\gamma_1(t) - 1)),$$

where $\gamma_1'(t) = \delta_1(t)$, $\gamma_1(t_0) = 1$, $\gamma_2(t_0) = 0$ and $\gamma_2'(t) = \delta_1(t)\delta_2(t)$. After applying an affine transformation we have shown:

Theorem 7. *Let M^3 be an indefinite proper affine hypersphere of $\mathbb{R}^4$ which admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry. Let $H = a_{22}^2 - a_4^2 (\neq 0)$ on M^3 . Then M^3 is affine equivalent with*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (t, v, w) \mapsto (\gamma_1(t)v, \gamma_1(t)w, \gamma_1(t)f(v, w) + \gamma_2(t), \gamma_1(t)),$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ is a positive definite improper affine sphere with affine normal $(0, 0, 1)$ and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve. Moreover, if M^3 admits a pointwise $SO(2)$ -symmetry then N^2 is an elliptic paraboloid.

We want to investigate the conditions imposed on the curve γ . For this we compute the derivatives of φ :

$$\begin{aligned} \varphi_t &= (\gamma_1' v, \gamma_1' w, \gamma_1' f(v, w) + \gamma_2', \gamma_1'), \\ \varphi_v &= (\gamma_1, 0, \gamma_1 f_v, 0), \quad \varphi_w = (0, \gamma_1, \gamma_1 f_w, 0), \\ \varphi_{tt} &= (\gamma_1'' v, \gamma_1'' w, \gamma_1'' f(v, w) + \gamma_2'', \gamma_1''), \\ \varphi_{tv} &= \frac{\gamma_1'}{\gamma_1} \varphi_v, \quad \varphi_{tw} = \frac{\gamma_1'}{\gamma_1} \varphi_w, \\ \varphi_{vv} &= (0, 0, f_{vv} \gamma_1, 0), \quad \varphi_{vw} = (0, 0, \gamma_1 f_{vw}, 0), \quad \varphi_{ww} = (0, 0, \gamma_1 f_{ww}, 0). \end{aligned} \tag{82}$$

M^3 is a proper hypersphere, i. e. $\xi = -H\varphi$. An easy computation shows that the condition that ξ is a transversal vector field, namely $0 \neq \det(\varphi_t, \varphi_v, \varphi_w, \xi) = -H\gamma_1^2(\gamma_1\gamma_2' - \gamma_1'\gamma_2)$, is equivalent to $\gamma_1 \neq 0$ and $\gamma_1\gamma_2' - \gamma_1'\gamma_2 \neq 0$. Since $(0, 0, 1, 0) = \frac{\gamma_1}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} \varphi_t - \frac{\gamma_1'}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} \varphi$, we have the following structure equations:

$$\begin{aligned} \varphi_{tt} &= \left(\frac{\gamma_1''}{\gamma_1} + \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{\gamma_1} \frac{\gamma_1}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} \right) \varphi_t + \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} \frac{1}{H} \xi, \\ \varphi_{tr} &= \frac{\gamma_1'}{\gamma_1} \varphi_r, \\ \varphi_{rs} &= \frac{\gamma_1^2}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} f_{rs} \varphi_t + \frac{\gamma_1\gamma_1'}{\gamma_1\gamma_2' - \gamma_1'\gamma_2} f_{rs} \frac{1}{H} \xi. \end{aligned} \tag{83}$$

We obtain that $\det h = h_{tt}(h_{vv}h_{ww} - h_{vw}^2) = \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{H^3(\gamma_1\gamma_2' - \gamma_1'\gamma_2)^3} \gamma_1^2(\gamma_1')^2(f_{vv}f_{ww} - f_{vw}^2)$. Since ψ is a positive definite improper affine sphere with affine normal $(0, 0, 1)$, we get by (4) that $f_{vv}f_{ww} - f_{vw}^2 = 1$. Now (4) (for ξ) is equivalent to $\gamma_1^4(\gamma_1\gamma_2' - \gamma_1'\gamma_2)^2 = \left| \frac{\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'}{(\gamma_1\gamma_2' - \gamma_1'\gamma_2)^3} \right| \gamma_1^2(\gamma_1')^2$. It follows that

$$\xi = -H\varphi \iff \gamma_1^2|\gamma_1\gamma_2' - \gamma_1'\gamma_2|^5 = |\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'|(\gamma_1')^2 \neq 0.$$

From the computations above also it follows that φ is indefinite iff either $\text{sign}(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2') = \text{sign}(H(\gamma_1\gamma_2' - \gamma_1'\gamma_2)) = -\text{sign}(\gamma_1\gamma_1')$ or $\text{sign}(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2') = -\text{sign}(H(\gamma_1\gamma_2' - \gamma_1'\gamma_2)) = -\text{sign}(\gamma_1\gamma_1')$.

Now we can formulate the converse theorem:

Theorem 8. *Let $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ be a positive definite improper affine sphere with affine normal $(0, 0, 1)$, and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve such that $\varphi(t, v, w) = (\gamma_1(t)v, \gamma_1(t)w, \gamma_1(t)f(v, w) + \gamma_2(t), \gamma_1(t))$ defines a 3-dimensional indefinite proper affine hypersphere. Then $\varphi(N^2 \times I)$ admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry.*

Moreover, if $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_1^2|\gamma_1\gamma_2' - \gamma_1'\gamma_2|^5 = -\text{sign}(\gamma_1\gamma_1')(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2')(\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.

Proof. We already have shown that φ defines a 3-dimensional indefinite proper affine hypersphere with affine normal $\xi = -H\varphi$. To prove the symmetry we need to compute K . We get the induced connection and the affine metric from the structure equations (83). We compute K using (6) and obtain:

$$\begin{aligned} (\nabla_{\varphi_t} h)(\varphi_r, \varphi_s) &= \left(\frac{\partial}{\partial t} \ln \left(\frac{\gamma_1 \gamma_1'}{\gamma_1 \gamma_2' - \gamma_1' \gamma_2} \right) - 2 \frac{\gamma_1'}{\gamma_1} \right) h(\varphi_r, \varphi_s), \\ (\nabla_{\varphi_r} h)(\varphi_t, \varphi_t) &= 0, \end{aligned}$$

implying that K_{φ_t} restricted to the space spanned by φ_v and φ_w is a multiple of the identity. Taking T in direction of φ_t , we see that φ_v and φ_w are orthogonal to T . Thus we can construct an ONB $\{T, V, W\}$ with V, W spanning $\text{span}\{\varphi_v, \varphi_w\}$ such that $a_1 = 2a_4$, $a_2 = a_3 = a_5 = 0$. By the considerations in Sec. 4 we see that φ admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry. $\square$

7.1.3 The third case: $\nu \equiv 0$ and $H = 0$ on M^3

The final cases now are that $\nu \equiv 0$ and $H = 0$ on the whole of M^3 and hence $a_{22} = \pm a_4$.

First we consider the case that $a_{22} = a_4 =: a > 0$. Again we use that M^3 admits a warped product structure and we fix a parameter t_0 . At the point t_0 , we have by (59)–(63):

$$\begin{aligned} D_V V &= +a_6 V - b_{23} W & +\xi, \\ D_V W &= +b_{23} V - a_6 W, \\ D_W V &= -(b_{33} + a_6) W, \\ D_W W &= +(b_{33} - a_6) V & +\xi, \\ D_X \xi &= 0. \end{aligned}$$

Thus, if v and w are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(t_0, v, w)$ as a positive definite improper affine sphere in a 3-dimensional linear subspace.

Moreover, we see that this improper affine sphere is a paraboloid provided that $a_6(t_0, v, w)$ vanishes identically. From the differential equations

(65) determining a_6 , we see that this is the case exactly when a_6 vanishes identically, i.e. when M^3 admits a pointwise $SO(2)$ -symmetry.

After applying an affine transformation and a change of coordinates, we may assume that

$$\varphi(t_0, v, w) = (v, w, f(v, w), 0), \quad (84)$$

with affine normal $\xi(t_0, v, w) = (0, 0, 1, 0)$, actually

$$\xi(t, v, w) = (0, 0, 1, 0)$$

(ξ is constant on M^3 by assumption). Furthermore we obtain by (57) and (58), that $D_U T = 2aU$ for all $U \in L_2$. We define $\delta := T - 2a\varphi$, which is transversal to $\text{span}\{V, W, \xi\}$. Since a is independent of v and w (cp. (64)), $D_U \delta = 0$, and we can assume that

$$T(t_0, v, w) - 2a(t_0)\varphi(t_0, v, w) = (0, 0, 0, 1). \quad (85)$$

We can integrate (64) ($T(a) = -4a^2$) and we take $a = \frac{1}{4t}$, $t > 0$. Thus (54) becomes $D_T T = -\frac{1}{2t}T - \xi$ and we obtain the following linear second order ordinary differential equation:

$$\frac{\partial^2}{\partial t^2} \varphi + \frac{1}{2t} \frac{\partial}{\partial t} \varphi = -\xi. \quad (86)$$

The general solution is $\varphi(t, v, w) = -\frac{t^2}{3}\xi + 2\sqrt{t}A(v, w) + B(v, w)$. The initial conditions (84) and (85) imply that $A(v, w) = (\frac{v}{2\sqrt{t_0}}, \frac{w}{2\sqrt{t_0}}, \frac{f(v, w)}{2\sqrt{t_0}} + \frac{2}{3}t_0^{3/2}, \sqrt{t_0})$ and $B(v, w) = (0, 0, -t_0^2, -2t_0)$. Obviously we can translate B to zero. Furthermore we can translate the affine sphere and apply an affine transformation to obtain $A(v, w) = \frac{1}{2\sqrt{t_0}}(v, w, f(v, w), 1)$. After a change of coordinates we get:

$$\varphi(t, v, w) = (tv, tw, tf(v, w) - ct^4, t), \quad c, t > 0. \quad (87)$$

Next we consider the case that $-a_{22} = a_4 =: a > 0$. Again we use that M^3 admits a warped product structure and we fix a parameter t_0 . A look at (59)–(63) suggests to define $\tilde{\xi} = -2aT + \xi$, then we get at the point t_0 :

$$\begin{aligned} D_V V &= +a_6 V - b_{23} W && +\tilde{\xi}, \\ D_V W &= +b_{23} V - a_6 W, \\ D_W V &= -(b_{33} + a_6)W, \\ D_W W &= +(b_{33} - a_6)V && +\tilde{\xi}, \\ D_U \tilde{\xi} &= 0. \end{aligned}$$

Thus, if v and w are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(t_0, v, w)$ as a positive definite improper affine sphere in a 3-dimensional linear subspace.

Moreover, we see that this improper affine sphere is a paraboloid provided that $a_6(t_0, v, w)$ vanishes identically. From the differential equations (65) determining a_6 , we see that this is the case exactly when a_6 vanishes identically, i.e. when M^3 admits a pointwise $SO(2)$ -symmetry.

After applying an affine transformation and a change of coordinates, we may assume that

$$\varphi(t_0, v, w) = (v, w, f(v, w), 0), \quad (88)$$

with affine normal

$$\tilde{\xi}(t_0, v, w) = (0, 0, 1, 0). \quad (89)$$

We have considered $\tilde{\xi}$ before. We can solve (70) ($\frac{\partial}{\partial t}\beta = -2a\beta$) explicitly by $\beta = c\frac{1}{\sqrt{a}}$ (cp. (64)) and get by (71)–(73) that $D_X(\frac{1}{\sqrt{a}}\tilde{\xi}) = 0$. Thus $\frac{1}{\sqrt{a}}(-2aT + \xi) =: C$ for a constant vector C , i. e. $T = -\frac{1}{2a}(\sqrt{a}C - \xi)$. Notice that by (63) ξ is a constant vector, too. We can choose $a = \frac{1}{4|t|}$, $t < 0$ (cp. (64)), and we obtain the ordinary differential equation:

$$\frac{\partial}{\partial t}\varphi = -\sqrt{|t|}C - 2t\xi, \quad t < 0. \quad (90)$$

The solution (after a translation) with respect to the initial condition (88) is $\varphi(t, v, w) = \frac{2}{3}|t|^{\frac{3}{2}}C - t^2\xi + (v, w, f(v, w), 0)$. Notice that C is a multiple of $\tilde{\xi}$ and hence by (89) a constant multiple of $(0, 0, 1, 0)$. Furthermore ξ is transversal to the space spanned by $\varphi(t_0, v, w)$. So we get after an affine transformation and a change of coordinates:

$$\varphi(t, v, w) = (v, w, f(v, w) + ct^3, t^4), \quad c, t > 0. \quad (91)$$

Combining both results (87) and (91) we have:

Theorem 9. *Let M^3 be an indefinite improper affine hypersphere of $\mathbb{R}^4$ which admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry. Let $a_{22}^2 = a_4^2$ on M^3 . Then M^3 is affine equivalent with either*

$$\begin{aligned} \varphi : I \times N^2 &\rightarrow \mathbb{R}^4 : (t, v, w) \mapsto (tv, tw, tf(v, w) - ct^4, t), \quad (a_{22} = a_4) \quad \text{or} \\ \varphi : I \times N^2 &\rightarrow \mathbb{R}^4 : (t, v, w) \mapsto (v, w, f(v, w) + ct^3, t^4), \quad (-a_{22} = a_4) \end{aligned}$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ is a positive definite improper affine sphere with affine normal $(0, 0, 1)$ and $c, t \in \mathbb{R}^+$.

Moreover, if M^3 admits a pointwise $SO(2)$ -symmetry then N^2 is an elliptic paraboloid.

The computations for the converse statement can be done completely analogous to the previous cases, they even are simpler (the curve is given parametrized).

Theorem 10. *Let $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ be a positive definite improper affine sphere with affine normal $(0, 0, 1)$. Define $\varphi(t, v, w) = (tv, tw, tf(v, w) - ct^4, t)$ or $\varphi(t, v, w) = (v, w, f(v, w) + ct^3, t^4)$, where $c, t \in \mathbb{R}^+$. Then φ defines a 3-dimensional indefinite improper affine hypersphere, which admits a pointwise $\mathbb{Z}_3$ - or $SO(2)$ -symmetry.*

7.2 Pointwise S_3 -symmetry

In the beginning of the section (cp. Lem. 10) we have shown that in the case of S_3 -symmetry, there exists a locally defined orthonormal frame $\{T, V, W\}$ such that K is given by:

$$\begin{aligned} K(T, T) &= 0, & K(T, V) &= 0, & K(T, W) &= 0, \\ K(V, V) &= a_6 V, & K(V, W) &= -a_6 W, & K(W, W) &= -a_6 W, \end{aligned}$$

(the vector field T (up to sign) and the function a_6 are invariantly defined) and the structure equations ((1), (2)) are:

$$D_T T = -\xi, \quad (92)$$

$$D_T V = +\frac{1}{3}a_{23}W, \quad (93)$$

$$D_T W = -\frac{1}{3}a_{23}V, \quad (94)$$

$$D_V T = +a_{22}V + a_{23}W, \quad (95)$$

$$D_W T = -a_{23}V + a_{22}W, \quad (96)$$

$$D_V V = +a_6 V - b_{23}W + a_{22}T + \xi, \quad (97)$$

$$D_V W = +b_{23}V - a_6 W + a_{23}T, \quad (98)$$

$$D_W V = - (b_{33} + a_6)W - a_{23}T, \quad (99)$$

$$D_W W = + (b_{33} - a_6)V + a_{22}T + \xi, \quad (100)$$

$$D_X \xi = -HX, \quad (101)$$

The Codazzi and Gauss equations ((12) and (13)) have the form (cp. Lem. 11 and 12):

$$T(\ln a_6) = -a_{22}, \quad V(\ln a_6) = 3b_{33}, \quad W(\ln a_6) = -3b_{23}, \quad (102)$$

$$\begin{aligned} T(a_{22}) &= -a_{22}^2 + a_{23}^2 + H, \\ V(a_{22}) &= W(a_{23}), \quad W(a_{22}) = -V(a_{23}), \end{aligned} \quad (103)$$

$$\begin{aligned} T(a_{23}) &= -2a_{22}a_{23}, \quad V(a_{23}) = -3(T(b_{23}) + a_{22}b_{23} + \frac{2}{3}a_{23}b_{33}), \\ W(a_{23}) &= 3(-T(b_{33}) + \frac{2}{3}a_{23}b_{23} - a_{22}b_{33}), \end{aligned} \quad (104)$$

$$V(b_{33}) - W(b_{23}) = -a_{22}^2 - \frac{5}{3}a_{23}^2 + b_{23}^2 + b_{33}^2 + H + 2a_6^2. \quad (105)$$

An evaluation of the equation $[V, W]a_6 = (\hat{\nabla}_V W - \hat{\nabla}_W V)a_6$ gives us in addition:

$$V(b_{23}) + W(b_{33}) = \frac{2}{3}a_{22}a_{23}. \quad (106)$$

Notice that by (102) and (103) the functions $|a_{22}|$ and $|a_{23}|$ are invariantly defined.

Positive definite affine hyperspheres admitting an S_3 -symmetry turn out to satisfy Chen's equality [Vra04]. They were studied in several articles ([SSVV97], [SV96], [KSV01], [KV99], [Vra04]) and now they are completely classified. Our structure and integrability conditions are kind of similar and luckily we can follow the approach in [KV99], solving the system of equations directly. We do expect that the approach of [KSV01] (reduction to a classification of 2-dimensional minimal surfaces in S_3^5 whose ellipses are circles, done for elliptic affine hyperspheres) also is generalizable to all cases.

From the structure equations (92) and (101) we get that $D_T D_T \varphi = H\varphi$ if $H = \pm 1$, resp. $D_T D_T \varphi = \text{const.}$, if $H = 0$. This implies for the integral curves γ of T that $\gamma'' = H\gamma$, thus (cp. [KSV01], Cor. 1):

Remark. φ is locally ruled by arcs of ellipses resp. hyperbolas if it is proper and $H = -1$ resp. $H = 1$, and by straight lines if it is improper ($H = 0$).

As a consequence of the ruling we can integrate the integrability conditions in T -direction explicitly. We introduce a coordinate function t by $\frac{\partial}{\partial t} := T$.

Lemma 21. *For any function t with $Tt = 1$ there is a dense open subset $\tilde{M} \subset M^3$ and there are functions $h, k, l: \tilde{M} \rightarrow \mathbb{R}$ with $Th = Tk = Tl = 0$ such that on $\tilde{M}$ we have if*

H = -1 then either $a_{22} = 0$, $a_{23} = \pm 1$, $a_6 = e^k$ or

$$a_{22} = -\frac{\sin(t+l)\cos(t+l)}{\cos^2(t+l) + \sinh^2(h)}, a_{23} = -\frac{\sinh(h)\cosh(h)}{\cos^2(t+l) + \sinh^2(h)},$$

$$a_6 = \frac{e^k}{\sqrt{\cos^2(t+l) + \sinh^2(h)}}.$$

H = 1 then either $a_{22} = \pm 1$, $a_{23} = 0$, $a_6 = e^{k \mp t}$ or

$$a_{22} = \frac{\sinh(t+l)\cosh(t+l)}{\cos^2(h) + \sinh^2(t+l)}, a_{23} = -\frac{\sin(h)\cos(h)}{\cos^2(h) + \sinh^2(t+l)},$$

$$a_6 = \frac{e^k}{\sqrt{\cos^2(h) + \sinh^2(t+l)}}.$$

H = 0 then either $a_{22} = 0$, $a_{23} = 0$, $a_6 = e^k$ or

$$a_{22} = \frac{t+l}{(t+l)^2 + h^2}, a_{23} = -\frac{h}{(t+l)^2 + h^2},$$

$$a_6 = \frac{e^k}{\sqrt{(t+l)^2 + h^2}}.$$

Proof. Equations (103) and (104) can be combined to $T(a_{22} + ia_{23}) = -(a_{22} + ia_{23})^2 + H$. Using the notation $y = -(a_{22} + ia_{23})$, we have to solve the first order differential equations $\frac{\partial}{\partial t}y = y^2 + 1$ ($H = -1$), $\frac{\partial}{\partial t}y = y^2 - 1$ ($H = 1$) resp. $\frac{\partial}{\partial t}y = y^2$ ($H = 0$). For $H = -1$ we get the generic solution $y = \tan(z + c)$ resp. the exceptional solution $y = \pm i$, for $H = 1$ we get $y = i \tan(z + c)$ resp. $y = \pm 1$ and for $H = 0$ we get $y = -\frac{1}{z}$ resp. $y = 0$. To obtain the real valued functions a_{22} and a_{23} along the ruling, we have to split $y(z(t)) = y(g(t) + ih(t))$ into real and imaginary part. In the first case ($H = -1$) we get (along the ruling) for example the generic solution: $y(t) = \tan(g(t) + ih(t))$ with $g(t) = t + l$ and $\frac{\partial}{\partial t}h = 0$. Now $y(t) = \frac{\sin(g(t)+ih)}{\cos(g(t)+ih)} = \frac{\sin(g(t))\cos(g(t))+i\sinh(h)\cosh(h)}{\cos^2(g(t))+\sinh^2(h)}$. Next we integrate (102) along the ruling to obtain a_6 . In the first case this gives generically: $-\ln a_6 = \frac{1}{2}\ln(\cos^2(t+l) + \sinh^2(h)) + \tilde{k}$. The other results are computed in the same way. $\square$

In order to solve the rest of the integrability conditions we need to introduce adapted coordinates. Since T is geometrically distinguished we would like to find Gaussian vector fields $\tilde{T}, \tilde{V}, \tilde{W}$ such that $\tilde{T} = T$. Analogous to [KV99] we look for functions c_1, c_2, b_1, b_2 such that

$$\begin{pmatrix} \tilde{V} \\ \tilde{W} \end{pmatrix} = \begin{pmatrix} c_1 & c_2 \\ -c_2 & c_1 \end{pmatrix} \left(\begin{pmatrix} V \\ W \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} T \right). \quad (107)$$

Lemma 22. *For $\tilde{T}, \tilde{V}, \tilde{W}$ to be (local) Gaussian vector fields it is necessary and sufficient that the following system of equations is satisfied:*

$$T(c_1) = a_{22}c_1 - \frac{2}{3}a_{23}c_2, \quad T(c_2) = \frac{2}{3}a_{23}c_1 + a_{22}c_2, \quad (108)$$

$$-V(c_1) + W(c_2) = b_{33}c_1 + b_{23}c_2, \quad V(c_2) + W(c_1) = b_{23}c_1 - b_{33}c_2, \quad (109)$$

$$T(b_1) = -a_{22}b_1 - \frac{2}{3}a_{23}b_2, \quad T(b_2) = \frac{2}{3}a_{23}b_1 - a_{22}b_2, \quad (110)$$

$$V(b_2) - W(b_1) = -2a_{23} + b_{23}b_1 + b_{33}b_2. \quad (111)$$

Proof. We have to show that our system of equations is equivalent to the vanishing of all commutators. Using $[\alpha X, \beta Y] = \alpha X(\beta)Y - \beta Y(\alpha)X + \alpha\beta(\widehat{\nabla}_X Y - \widehat{\nabla}_Y X)$ we get

$$\begin{aligned} [V, W] &= b_{23}V + b_{33}W + 2a_{23}T, & [T, V] &= -a_{22}V - \frac{2}{3}a_{23}W, \\ [T, W] &= \frac{2}{3}a_{23}V - a_{22}W. \end{aligned} \quad (112)$$

Therefore

$$\begin{aligned} [\tilde{T}, \tilde{V}] &= [T, c_1V + c_2W + (c_1b_1 + c_2b_2)T] \\ &= (T(c_1) - a_{22}c_1 + \frac{2}{3}a_{23}c_2)V + (T(c_2) - \frac{2}{3}a_{23}c_1 - a_{22}c_2)W \\ &\quad + (T(c_1)b_1 + c_1T(b_1) + T(c_2)b_2 + c_2T(b_2))T. \end{aligned}$$

Hence $[\tilde{T}, \tilde{V}] = 0$ is equivalent to (108) and the equation

$$c_1 T(b_1) + c_2 T(b_2) = c_1(-a_{22}b_1 - \frac{2}{3}a_{23}b_2) + c_2(\frac{2}{3}a_{23}b_1 - a_{22}b_2). \quad (113)$$

Similarly we get

$$\begin{aligned} [\tilde{T}, \tilde{W}] &= [T, -c_2V + c_1W + (-c_2b_1 + c_1b_2)T] \\ &= (-T(c_2) + a_{22}c_2 + \frac{2}{3}a_{23}c_1)V + (T(c_1) + \frac{2}{3}a_{23}c_2 - a_{22}c_1)W \\ &\quad + (-T(c_2)b_1 - c_2T(b_1) + T(c_1)b_2 + c_1T(b_2))T. \end{aligned}$$

Hence $[\tilde{T}, \tilde{W}] = 0$ is equivalent to (108) and the equation

$$-c_2T(b_1) + c_1T(b_2) = -c_2(-a_{22}b_1 - \frac{2}{3}a_{23}b_2) + c_1(\frac{2}{3}a_{23}b_1 - a_{22}b_2). \quad (114)$$

Since $c_1^2 + c_2^2 \neq 0$, equations (113) and (114) are equivalent to (110). Finally, a lengthy but straightforward calculation gives (we use the abbreviation $\alpha = c_1b_1 + c_2b_2$ and $\beta = -c_2b_1 + c_1b_2$)

$$\begin{aligned} [\tilde{V}, \tilde{W}] &= [c_1V + c_2W + \alpha T, -c_2V + c_1W + \beta T] \\ &= \{c_1(-V(c_2) - W(c_1)) + c_2(V(c_1) - W(c_2)) + (c_1^2 + c_2^2)b_{23} \\ &\quad - \alpha(T(c_2) - c_2a_{22} - c_1\frac{2}{3}a_{23}) - \beta(T(c_1) - c_1a_{22} + c_2\frac{2}{3}a_{23})\}V \\ &\quad + \{-c_2(-V(c_2) - W(c_1)) + c_1(V(c_1) - W(c_2)) + (c_1^2 + c_2^2)b_{33} \\ &\quad + \alpha(T(c_1) - c_1a_{22} + c_2\frac{2}{3}a_{23}) - \beta(T(c_2) - c_2a_{22} - c_1\frac{2}{3}a_{23})\}W \\ &\quad + \{c_1(V(\beta) - W\alpha) + c_2(V\alpha + W(\beta)) + (c_1^2 + c_2^2)2a_{23} \\ &\quad + \alpha T(\beta) - \beta T(\alpha)\}T. \end{aligned}$$

Using equation (108), the vanishing of the V- and W-component of $[\tilde{V}, \tilde{W}]$ is equivalent to (109). Now $[\tilde{V}, \tilde{W}]$ vanishes in addition to $[\tilde{T}, \tilde{V}]$ and $[\tilde{T}, \tilde{W}]$ if and only if its V-, W-components vanish and the equation

$$0 = (c_1^2 + c_2^2)(2a_{23} - b_1b_{23} - b_2b_{33} + V(b_2) - W(b_1))$$

holds. However, $c_1^2 + c_2^2 \neq 0$, thus the last equation is equivalent to (111). $\square$

To obtain adapted coordinates we need to solve the system of equations (108)–(111). First we will treat the generic cases.

Lemma 23. *Assume that we are in one of the generic cases (cp. Lem. 21) and let f be any function with $Tf = -1$. Then the functions c_1, c_2, b_1, b_2 , defined by*

$$(c_1 + ic_2)^3 = \frac{1}{a_6((a_{22} + ia_{23})^2 - H)}, \quad b_1 = V(f), \quad b_2 = W(f), \quad (115)$$

are a solution of the system of differential equations (108)–(111).

Proof. Observe first that the system of equations splits into two independent subsystems, namely the system (108), (109), and the system (110), (111). Assume first that da_{22} , da_{23} and da_6 are everywhere linearly independent. We start with the system (108), (109). We reformulate the system with respect to the coordinate functions a_{22} , a_{23} , $a = \ln a_6$, using the integrability conditions (102)–(104):

$$\begin{aligned} a_{22}c_1 - \frac{2}{3}a_{23}c_2 &= T(c_1) \\ &= ((-a_{22}^2 + a_{23}^2 + H)\frac{\partial}{\partial a_{22}} - 2a_{22}a_{23}\frac{\partial}{\partial a_{23}} - a_{22}\frac{\partial}{\partial a})c_1, \end{aligned} \quad (116)$$

$$\begin{aligned} \frac{2}{3}a_{23}c_1 + a_{22}c_2 &= T(c_2) \\ &= (-a_{22}^2 + a_{23}^2 + H)\frac{\partial}{\partial a_{22}} - 2a_{22}a_{23}\frac{\partial}{\partial a_{23}} - a_{22}\frac{\partial}{\partial a})c_2, \end{aligned} \quad (117)$$

$$\begin{aligned} b_{33}c_1 + b_{23}c_2 &= -V(c_1) + W(c_2) \\ &= V(a_{22})\left(-\frac{\partial}{\partial a_{22}}c_1 + \frac{\partial}{\partial a_{23}}c_2\right) + W(a_{22})\left(\frac{\partial}{\partial a_{22}}c_2 + \frac{\partial}{\partial a_{23}}c_1\right) \\ &\quad - 3b_{33}\frac{\partial}{\partial a}c_1 - 3b_{23}\frac{\partial}{\partial a}c_2, \end{aligned} \quad (118)$$

$$\begin{aligned} b_{23}c_1 - b_{33}c_2 &= V(c_2) + W(c_1) \\ &= V(a_{22})\left(\frac{\partial}{\partial a_{22}}c_2 + \frac{\partial}{\partial a_{23}}c_1\right) + W(a_{22})\left(\frac{\partial}{\partial a_{22}}c_1 - \frac{\partial}{\partial a_{23}}c_2\right) \\ &\quad + 3b_{33}\frac{\partial}{\partial a}c_2 - 3b_{23}\frac{\partial}{\partial a}c_1. \end{aligned} \quad (119)$$

The last two equations are satisfied if (c_1, c_2) satisfy the Cauchy-Riemann equations with respect to the coordinates a_{22} and a_{23} for any fixed a , i. e.

$$\frac{\partial}{\partial a_{22}}c_1 = \frac{\partial}{\partial a_{23}}c_2, \quad \frac{\partial}{\partial a_{22}}c_2 = -\frac{\partial}{\partial a_{23}}c_1$$

and their a -dependence is given by

$$\frac{\partial}{\partial a}c_1 = -\frac{1}{3}c_1, \quad \frac{\partial}{\partial a}c_2 = -\frac{1}{3}c_2. \quad (120)$$

Therefore we assume now that $c_1(z) + ic_2(z)$ is a holomorphic function with respect to $z = a_{22} + ia_{23}$ and that (120) holds. Since $(z^2 - H)\frac{\partial}{\partial z}(c_1 + ic_2) = ((a_{22}^2 - a_{23}^2 - H)\frac{\partial}{\partial a_{22}} + 2a_{22}a_{23}\frac{\partial}{\partial a_{23}})c_1 + i((+a_{22}^2 - a_{23}^2 - H)\frac{\partial}{\partial a_{22}} + 2a_{22}a_{23}\frac{\partial}{\partial a_{23}})c_2$ and $-\frac{2}{3}z(c_1 + ic_2) = -\frac{2}{3}(a_{22}c_1 - a_{23}c_2 + i(a_{22}c_2 + a_{23}c_1))$, equations (116) and (117) are equivalent to

$$(z^2 - H)\frac{\partial}{\partial z}(c_1 + ic_2) = -\frac{2}{3}z(c_1 + ic_2).$$

We can integrate the last equation and obtain that $(c_1 + ic_2)^{-3} = g(a)((a_{22} + ia_{23})^2 - H)$. Finally equation (120) implies $g(a) = e^a = a_6$. The second

subsystem can be rewritten as the integrability conditions for a function f with $Tf = -1$. In fact, if we set $b_1 = V(f)$, $b_2 = W(f)$ and $b_3 = T(f)$, we get by (110) and (111):

$$\begin{aligned} [V, T](f) &= a_{22}V(f) + \frac{2}{3}a_{23}W(f) = a_{22}b_1 + \frac{2}{3}a_{23}b_2 = -T(b_1), \\ [W, T](f) &= -\frac{2}{3}a_{23}V(f) + a_{22}W(f) = -\frac{2}{3}a_{23}b_1 + a_{22}b_2 = -T(b_2), \\ [V, W](f) &= 2a_{23}b_3 + b_{23}b_1 + b_{33}b_2 = 2a_{23}(b_3 + 1) + V(b_2) - W(b_1). \end{aligned}$$

Hence these integrability conditions are satisfied for $b_3 = -1$ and arbitrary functions b_1 and b_2 .

Now we deal with the case that da_{22} , da_{23} and da_6 are linearly dependent in some subset of M^3 . We can still define functions c_1, c_2, b_1, b_2 by equations (115). A direct computation shows that these functions satisfy the system of equations (108)–(111). $\square$

We denote the coordinates provided by Lemma 23 by (t, v, w) , i. e. $\frac{\partial}{\partial t} = \tilde{T} = T$, $\frac{\partial}{\partial v} = \tilde{V}$ and $\frac{\partial}{\partial w} = \tilde{W}$. We will now make a special choice for f in order to simplify equations in Lemma 21, namely to make $l = 0$.

Lemma 24. *There exist a function f with $T(f) = -1$ and a constant c such that after replacing t by $t + c$ one has $l = 0$ in Lemma 21. The function f is given by*

$$\mathbf{H} = -\mathbf{1}: f = -\frac{1}{2} \arcsin \left(\frac{2a}{\sqrt{(-a_{22}^2 + a_{23}^2 - 1)^2 + 4a_{22}^2 a_{23}^2}} \right),$$

$$\mathbf{H} = \mathbf{1}: f = -\frac{1}{2} \operatorname{arcsinh} \left(\frac{2a}{\sqrt{(-a_{22}^2 + a_{23}^2 + 1)^2 + 4a_{22}^2 a_{23}^2}} \right),$$

$$\mathbf{H} = \mathbf{0}: f = -\frac{a}{\sqrt{(-a_{22}^2 + a_{23}^2)^2 + 4a_{22}^2 a_{23}^2}}.$$

The variable t is given by $t = -f + \text{const.}$

Proof. A direct calculation shows that our choice of f satisfies $T(f) = -1$. Since $V(f) = b_1$, $W(f) = b_2$, we get from (107) that

$$\begin{pmatrix} \frac{\partial}{\partial v} \\ \frac{\partial}{\partial w} \end{pmatrix} f = \begin{pmatrix} c_1 & c_2 \\ -c_2 & c_1 \end{pmatrix} \left(\begin{pmatrix} V(f) \\ W(f) \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} T(f) \right) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Using Lemma 21 we compute that $f = -(t + l)$. The function l must be a constant, since $Tl = 0$ by Lemma 21 and we have just shown that also $\frac{\partial}{\partial v} f = \frac{\partial}{\partial w} f = 0$. $\square$

In the generic cases we have everything to prove the final results:

Theorem 11. Let M^3 be an indefinite affine hypersphere of $\mathbb{R}^4$ with $H = -1$ which admits a pointwise S_3 -symmetry and assume, that the invariant functions everywhere satisfy $(a_{22}, a_{23}) \neq (0, \pm 1)$. Then M^3 admits coordinates (t, v, w) and there are functions $h, k: (v, w) \rightarrow \mathbb{R}$ such that

$$\begin{aligned} a_{22} &= -\frac{\sin(t) \cos(t)}{\cos^2(t) + \sinh^2(h)}, & a_{23} &= -\frac{\sinh(h) \cosh(h)}{\cos^2(t) + \sinh^2(h)}, \\ a_6 &= \frac{e^k}{\sqrt{\cos^2(t) + \sinh^2(h)}}, \end{aligned}$$

where h, k satisfy the system of differential equations

$$\left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)h = e^{-\frac{2}{3}k} \sinh(2h), \quad \left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)k = 3e^{-\frac{2}{3}k}(2e^{2k} - \cosh(2h)). \quad (121)$$

Theorem 12. Let M^3 be an indefinite affine hypersphere of $\mathbb{R}^4$ with $H = 1$ which admits a pointwise S_3 -symmetry and assume, that the invariant functions everywhere satisfy $(a_{22}, a_{23}) \neq (\pm 1, 0)$. Then M^3 admits coordinates (t, v, w) and there are functions $h, k: (v, w) \rightarrow \mathbb{R}$ such that

$$\begin{aligned} a_{22} &= \frac{\sinh(t) \cosh(t)}{\cos^2(h) + \sinh^2(t)}, & a_{23} &= -\frac{\sin(h) \cos(h)}{\cos^2(h) + \sinh^2(t)}, \\ a_6 &= \frac{e^k}{\sqrt{\cos^2(h) + \sinh^2(t)}}, \end{aligned}$$

where h, k satisfy the system of differential equations

$$\left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)h = -e^{-\frac{2}{3}k} \sin(2h), \quad \left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)k = 3e^{-\frac{2}{3}k}(2e^{2k} + \cos(2h)). \quad (122)$$

Theorem 13. Let M^3 be an indefinite improper affine hypersphere of $\mathbb{R}^4$ which admits a pointwise S_3 -symmetry and assume, that the invariant functions everywhere satisfy $(a_{22}, a_{23}) \neq (0, 0)$. Then M^3 admits coordinates (t, v, w) and there are functions $h, k: (v, w) \rightarrow \mathbb{R}$ such that

$$a_{22} = \frac{t}{t^2 + h^2}, \quad a_{23} = -\frac{h}{t^2 + h^2}, \quad a_6 = \frac{e^k}{\sqrt{t^2 + h^2}},$$

where h, k satisfy the system of differential equations

$$\left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)h = 2e^{-\frac{2}{3}k}h, \quad \left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)k = 3e^{-\frac{2}{3}k}(2e^{2k} - 1). \quad (123)$$

Proof of the theorems. Let M^3 be an indefinite affine hypersphere of $\mathbb{R}^4$ which admits a pointwise S_3 -symmetry and assume, that the invariant functions everywhere satisfy $(a_{22}, a_{23}) \neq (0, \pm 1)$ ($H = -1$), $(a_{22}, a_{23}) \neq (\pm 1, 0)$ ($H = 1$) resp. $(a_{22}, a_{23}) \neq (0, 0)$ ($H = 0$). By Lemma 21 we can assume without loss of generality that the functions a_{22} , a_{23} and a_6 are given by the equations above. We need to show that all integrability conditions ((102)–(106)) are satisfied. To simplify the computations we will work in some situations in $\mathbb{C}$ and use the identification

$$x + iy \longleftrightarrow \begin{pmatrix} x & -y \\ y & x \end{pmatrix}$$

The equations (103) then are equivalent to

$$(V + iW)(a_{23}) = i(V + iW)(a_{22}) \quad (124)$$

(and $T(a_{22}) = -a_{22}^2 + a_{23}^2 + H$). To get $V(a_{22})$, $V(a_{23})$, $W(a_{22})$ and $W(a_{23})$, first we use (107) and the notation $c = c_1 + ic_2$, $b = b_1 + ib_2$, $\partial = \frac{\partial}{\partial v} + i\frac{\partial}{\partial w}$:

$$(V + iW) = \frac{1}{c}\partial - bT. \quad (125)$$

For the computation we would like to know ∂a_{22} and ∂a_{23} resp. $X(a_{22})$ and $X(a_{23})$ for $X \in \{\frac{\partial}{\partial v}, \frac{\partial}{\partial w}\}$. In the proof of Lemma 21 we have used the notation $y = -(a_{22} + ia_{23})$, and we have shown that $y = \tan(z + c) = \tan(t + ih)$ ($H = -1$), $y = i \tan(z + c) = i \tan(h + it)$ ($H = 1$) resp. $y = -\frac{1}{z} = -\frac{1}{t + ih}$ ($H = 0$), where $T(h) = 0$. Now $\frac{\partial y}{\partial z} \frac{\partial z}{\partial t} = T(y) = -(T(a_{22}) + iT(a_{23}))$, i. e. $\frac{\partial y}{\partial z} = -(T(a_{22}) + iT(a_{23}))$ ($H = -1, 0$) resp. $\frac{\partial y}{\partial z} = -T(a_{23}) + iT(a_{22})$ ($H = 1$), and furthermore $-(X(a_{22}) + iX(a_{23})) = X(y) = \frac{\partial y}{\partial z} X(z)$ with $X(z) = iX(h)$ ($H = -1, 0$) resp. $X(z) = X(h)$ ($H = 1$). Thus we get with $\varepsilon_H = \begin{cases} 1 & \text{if } H = -1, 0, \\ -1 & \text{if } H = 1, \end{cases}$

$$X(a_{22}) = -\varepsilon_H T(a_{23})X(h), \quad X(a_{23}) = \varepsilon_H T(a_{22})X(h), \quad (126)$$

Applying (125) and (126) to (124), we obtain that $\frac{1}{c}\varepsilon_H T(a_{22})\partial(h) - bT(a_{23}) = (V + iW)(a_{23}) = i(V + iW)(a_{22}) = -\frac{1}{c}i\varepsilon_H T(a_{23})\partial(h) - ibT(a_{22})$. Thus equation (124) is equivalent to

$$b = \varepsilon_H \frac{1}{c} i \partial(h). \quad (127)$$

Now equations (102) are equivalent to:

$$\begin{aligned} 3(b_{33} - ib_{23}) &= (V + iW)(\ln a_6) = \frac{1}{c}\partial(\ln a_6) - b(-a_{22}) \\ &= \frac{1}{c}(\partial(\ln a_6) + \varepsilon_H i \partial(h)a_{22}). \end{aligned} \quad (128)$$

In complex notion the integrability conditions (104) are

$$(V + iW)(a_{23}) = -(3iT + 3ia_{22} + 2a_{23})(b_{33} - ib_{23}). \quad (129)$$

From the above we already know that

$$(V + iW)(a_{23}) = \frac{1}{\bar{c}} \varepsilon_H \partial(h) (T(a_{22}) - iT(a_{23})). \quad (130)$$

For the right side of equation (129) we need to know $T(\bar{c})$. Using (108) we obtain that $T(\bar{c}) = -\frac{1}{3\bar{c}}(3a_{22} - ia_{23})$. With the help of (126) and the equations for $T(a_{22})$ and $T(a_{23})$ (cp. (103) and (104)) we get:

$$\begin{aligned} 3T(b_{33} - ib_{23}) &= \frac{1}{3\bar{c}} \{ \partial(\ln a_6)(-3a_{22} + ia_{23}) \\ &\quad + \varepsilon_H \partial(h)(ia_{22}(-3a_{22} + ia_{23}) + 3T(a_{23}) + i3T(a_{22})) \} \\ &= \frac{1}{3\bar{c}} \{ \partial(\ln a_6)(-3a_{22} + ia_{23}) \\ &\quad + \varepsilon_H \partial(h)(-4a_{22}a_{23} + i(-6a_{22}^2 + 3a_{23}^2 + 3H)) \}. \end{aligned} \quad (131)$$

From (130) and (131) (and (128)) it is easy to check that (129) is true and thus (104). For the investigation of (105) and (106) we first compute the left hand sides in complex notion. We use (125), the notion $\Delta = \frac{\partial^2}{\partial^2 v} + \frac{\partial^2}{\partial^2 w}$, and also that by (128) $b_{33} + ib_{23} = \frac{1}{3c}(\overline{\partial(\ln a_6)} - ia_{22}\varepsilon_H \overline{\partial(h)})$, and obtain after a while:

$$\begin{aligned} (V + iW)(b_{33} + ib_{23}) &= \frac{1}{3i|c|^2} \{ (a_{22}\varepsilon_H \overline{\partial(h)} + i\overline{\partial(\ln a_6)})(-\frac{1}{c})(\partial c - \varepsilon_H i\partial(h)T(c)) \\ &\quad + \varepsilon_H a_{22}\Delta h + i\Delta(\ln a_6) - iT(a_{22})|\partial(h)|^2 \}. \end{aligned} \quad (132)$$

The complex function c was defined by $c = (e^{\ln a_6}(z^2 - H))^{-\frac{1}{3}}$ (cp. (115)), where $z = a_{22} + ia_{23}$. Thus $\partial c = -\frac{c}{3}(\partial(\ln a_6) + \frac{2z\varepsilon_H \partial(z)\partial(h)}{z^2 - H}) = -\frac{c}{3}(\partial(\ln a_6) - 2\varepsilon_H i(a_{22} + ia_{23})\partial(h))$ (cp. (126)). By (108) we have that $T(c) = (a_{22} + i\frac{2}{3}a_{23})c$. So we can simplify $\partial c - \varepsilon_H i\partial(h)T(c) = -\frac{c}{3}(\partial(\ln a_6) + \varepsilon_H a_{22}i\partial(h))$, and hence (132) becomes:

$$\begin{aligned} (V + iW)(b_{33} + ib_{23}) &= \frac{1}{3|c|^2} \{ -ia_{22}\varepsilon_H \frac{1}{3} 2i\Im(\overline{\partial(h)}\partial(\ln a_6)) + \frac{1}{3}|\partial(\ln a_6)|^2 \\ &\quad + |\partial(h)|^2(\frac{4}{3}a_{22}^2 - a_{23}^2 - H) + \Delta(\ln a_6) - i\varepsilon_H a_{22}\Delta h \}. \end{aligned} \quad (133)$$

Splitting into real and imaginary part we have:

$$\begin{aligned} V(b_{33}) - W(b_{23}) = & \frac{1}{3|c|^2} \{ 2a_{22}\varepsilon_H \frac{1}{3} \Im(\overline{\partial(h)} \partial(\ln a_6)) + \frac{1}{3} |\partial(\ln a_6)|^2 \\ & + |\partial(h)|^2 (\frac{4}{3} a_{22}^2 - a_{23}^2 - H) + \Delta(\ln a_6) \}, \end{aligned} \quad (134)$$

$$V(b_{23}) + W(b_{33}) = -\frac{1}{3|c|^2} \varepsilon_H a_{22} \Delta h. \quad (135)$$

This means that (106) is equivalent to

$$\Delta h = -2\varepsilon_H |c|^2 b. \quad (136)$$

Furthermore by (128) we get $b_{33}^2 + b_{23}^2 = |b_{33} + ib_{23}|^2 = \frac{1}{9|c|^2} (|\partial(\ln a_6)|^2 + a_{22}|\partial(h)|^2 + 2\varepsilon_H a_{22} \Im(\overline{\partial(h)} \partial(\ln a_6)))$ and therefore:

$$V(b_{33}) - W(b_{23}) - (b_{33}^2 + b_{23}^2) = \frac{1}{3|c|^2} (|\partial(h)|^2 (a_{22}^2 - a_{23}^2 - H) + \Delta(\ln a_6)). \quad (137)$$

To simplify the right hand side we need to compute $\Delta(\ln a_6)$. A lengthy but straightforward computation, using the definition of a_6 in each case, leads to:

$$V(b_{33}) - W(b_{23}) - (b_{33}^2 + b_{23}^2) = \frac{1}{3|c|^2} (\Delta k + \varepsilon_H a_{23} \Delta h). \quad (138)$$

If we use (136), we get from (138) that (105) is equivalent to

$$\Delta k = 3|c|^2 (-a_{22}^2 - a_{23}^2 + H + 2a_6^2). \quad (139)$$

Finally, to prove the equivalence of (136) and (139) with the systems of differential equations (cp. (121), (122), (123)), one needs to juggle with trigonometric equalities. Explicitly we will give one more step:

$$\mathbf{H} = -\mathbf{1} : |c|^2 = e^{-\frac{2}{3}k} (\cos^2(t) + \sinh^2(h)),$$

$$\mathbf{H} = \mathbf{1} : |c|^2 = e^{-\frac{2}{3}k} (\cos^2(h) + \sinh^2(t)),$$

$$\mathbf{H} = \mathbf{0} : |c|^2 = e^{-\frac{2}{3}k} (h^2 + t^2).$$

□

We are left with the exceptional cases (cp. Lemma 21). Using (112) it is easy to show that the Gauss equations (104) and (106) are just the integrability conditions for a function k with

$$T(k) = 0, \quad V(k) = 3b_{33}, \quad W(k) = -3b_{23}. \quad (140)$$

For $a_6 = \exp(-a_{22}t+k)$ (cp. Lemma 21) the latter equations are the Codazzi equations (102). To solve the last non-trivial integrability condition (105) we need to find adapted coordinates, i. e. by Lemma 22 we need to solve the system of equations (108)–(111). Notice that for $H = 0$ or $H = 1$ we have $a_{23} = 0$, hence by (112) the distribution $T^\perp = \text{span}\{V, W\}$ is integrable and we do not need the functions b_1, b_2 in (107).

Lemma 25. *Assume that we are in one of the exeptional cases (cp. Lem. 21) and let f be any function with $Tf = -1$. Then the functions c_1, c_2, b_1, b_2 , defined by*

$$(c_1 + ic_2)^3 = \frac{1}{a_6} e^{2(a_{22} + ia_{23})t}, \quad b_1 = V(f), \quad b_2 = W(f), \quad (141)$$

are a solution of the system of differential equations (108)–(111).

Proof. Observe first that the system of equations splits into two independent subsystems, namely the system (108), (109), and the system (110), (111). We rewrite the first systems, using again the complex notation $c = c_1 + ic_2$: (108) is equivalent to $T(c) = (a_{22} + i\frac{2}{3}a_{23})c$ and (109) to

$$(V + iW)(c) = (-b_{33} + ib_{23})c = -\frac{1}{3}(V + iW)(k)c. \quad (142)$$

Integrating these equations we get that $c = \exp(-\frac{1}{3}k + (a_{22} + i\frac{2}{3}a_{23})t)$. Since $a_6 = \exp(-a_{22}t + k)$, we get the expression for c . The proof for the second subsystem (only non-trivial for $H = -1$) is the same as before (cp. Lemma 23). $\square$

As before we denote the coordinates, this time provided by Lemma 25, by (t, v, w) , i. e. $\frac{\partial}{\partial t} = \tilde{T} = T$, $\frac{\partial}{\partial v} = \tilde{V}$ and $\frac{\partial}{\partial w} = \tilde{W}$. Now we are ready to prove the final results for the exeptional cases:

Theorem 14. *Let M^3 be an indefinite affine hypersphere of $\mathbb{R}^4$ with $H = -1$ which admits a pointwise S_3 -symmetry and assume, that the invariant functions everywhere satisfy $(a_{22}, a_{23}) = (0, \pm 1)$. Then M^3 admits coordinates (t, v, w) and there is a function $k: (v, w) \rightarrow \mathbb{R}$ such that $a_6 = e^k$, where k satisfies the elliptic pde*

$$\left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)k = 2e^{-\frac{2}{3}k}(3e^{2k} - 4). \quad (143)$$

Theorem 15. *Let M^3 be an indefinite affine hypersphere of $\mathbb{R}^4$ with $H = 1$ which admits a pointwise S_3 -symmetry and assume, that the invariant functions everywhere satisfy $(a_{22}, a_{23}) = (\pm 1, 0)$. Then M^3 admits coordinates (t, v, w) and there is a function $k: (v, w) \rightarrow \mathbb{R}$ such that $a_6 = e^{k \mp t}$, where k satisfies the elliptic pde*

$$\left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)k = 6e^{\frac{4}{3}k}. \quad (144)$$

Theorem 16. *Let M^3 be an indefinite improper affine hypersphere of $\mathbb{R}^4$ which admits a pointwise S_3 -symmetry and assume, that the invariant functions everywhere satisfy $(a_{22}, a_{23}) = (0, 0)$. Then M^3 admits coordinates (t, v, w) and there is a function $k: (v, w) \rightarrow \mathbb{R}$ such that $a_6 = e^k$, where k satisfies the elliptic pde*

$$\left(\frac{\partial^2}{\partial v^2} + \frac{\partial^2}{\partial w^2}\right)k = 6e^{\frac{4}{3}k}. \quad (145)$$

Proof of the theorems. We already have seen that the integrability conditions (102)–(104) and (106) are satisfied. Hence we need to show that equation (105) is true. Using (106) and (140) we get in complex notation $V(b_{33}) - W(b_{23}) = (V + iW)(b_{33} + ib_{23}) = (V + iW)\frac{1}{3}(V + iW)(k)$. We have found coordinates such that $V + iW = \frac{c}{|c|^2}\partial - bT$ (cp. (107)), here $\partial = \frac{\partial}{\partial v} + i\frac{\partial}{\partial w} = \tilde{V} + i\tilde{W}$ and $b = b_1 + ib_2$. Thus $(V + iW)\frac{1}{3}(V + iW)(k) = \frac{1}{3}(V + iW)(\frac{c}{|c|^2}\partial(k) - \bar{b}T(k)) = \frac{1}{3}(V + iW)(\frac{1}{c}\partial(k))$, since $T(k) = 0$ by (140). We know $(V + iW)(c)$ by equation (142), and because of $T(\partial(k)) = 0$ we finally obtain that

$$V(b_{33}) - W(b_{23}) = \frac{1}{9|c|^2}(|\partial(k)|^2 + 3\Delta k). \quad (146)$$

Also from (142) follows that $b_{22}^2 + b_{23}^2 = \frac{1}{9}|(V + iW)(k)|^2 = \frac{1}{9|c|^2}|\partial(k)|^2$, thus (105) is equivalent to

$$\frac{1}{3|c|^2}\Delta k = H - a_{22}^2 - \frac{5}{3}a_{23}^2 + 2a_6^2.$$

To complete the proof we only need to replace H , a_{22} , a_{23} , a_6 and c in each case according to their definition. $\square$

8 Pointwise $SO(1, 1)$ -symmetry

Let M^3 be a hypersphere admitting a $SO(1, 1)$ -symmetry. According to Thm. 2, there exists for every $p \in M^3$ an LVB $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ of $T_p M^3$ such that

$$\begin{aligned} K(\mathbf{v}, \mathbf{v}) &= -2b_4\mathbf{v}, & K(\mathbf{v}, \mathbf{e}) &= b_4\mathbf{e}, & K(\mathbf{v}, \mathbf{f}) &= b_4\mathbf{f}, \\ K(\mathbf{e}, \mathbf{e}) &= 0, & K(\mathbf{e}, \mathbf{f}) &= b_4\mathbf{v}, & K(\mathbf{f}, \mathbf{f}) &= 0, \end{aligned}$$

where $b_4 > 0$.

We would like to extend the LVB locally. It is well known that $\widehat{\text{Ric}}$ (cp. (7)) is a symmetric operator and we compute (some of the computations in this section are done with the CAS Mathematica⁴):

⁴http://www.math.tu-berlin.de/~schar/IndefSym_typ8.html

Lemma 26. *Let $p \in M^3$ and $\{\mathbf{e}, \mathbf{v}, \mathbf{f}\}$ the basis constructed earlier. Then*

$$\begin{aligned}\widehat{\text{Ric}}(\mathbf{v}, \mathbf{v}) &= 2(H + 3b_4^2), & \widehat{\text{Ric}}(\mathbf{v}, \mathbf{e}) &= 0, \\ \widehat{\text{Ric}}(\mathbf{v}, \mathbf{f}) &= 0, & \widehat{\text{Ric}}(\mathbf{e}, \mathbf{e}) &= 0, \\ \widehat{\text{Ric}}(\mathbf{e}, \mathbf{f}) &= 2(H + b_4^2), & \widehat{\text{Ric}}(\mathbf{w}, \mathbf{w}) &= 0.\end{aligned}$$

Proof. The proof is a straight-forward computation using the Gauss equation (13). It follows e. g. that

$$\begin{aligned}\hat{R}(\mathbf{v}, \mathbf{e})\mathbf{v} &= -H\mathbf{e} - K_{\mathbf{v}}(b_4\mathbf{e}) + K_{\mathbf{e}}(-2b_4\mathbf{v}) = -H\mathbf{e} - b_4^2\mathbf{e} - 2b_4^2\mathbf{e} \\ &= -(H + 3b_4^2)\mathbf{e}, \\ \hat{R}(\mathbf{v}, \mathbf{f})\mathbf{v} &= -H\mathbf{f} - K_{\mathbf{v}}(b_4\mathbf{f}) + K_{\mathbf{f}}(-2b_4\mathbf{v}) = -H\mathbf{f} - b_4^2\mathbf{f} - 2b_4^2\mathbf{f} \\ &= -(H + 3b_4^2)\mathbf{f}, \\ \hat{R}(\mathbf{f}, \mathbf{e})\mathbf{v} &= -K_{\mathbf{f}}(b_4\mathbf{e}) + K_{\mathbf{e}}(b_4\mathbf{f}) = 0.\end{aligned}$$

From this it immediately follows that

$$\widehat{\text{Ric}}(\mathbf{v}, \mathbf{v}) = 2(H + 3b_4^2)$$

and

$$\widehat{\text{Ric}}(\mathbf{v}, \mathbf{e}) = 0.$$

The other equations follow by similar computations. $\square$

Remark. For the scalar curvature of M^3 we obtain $\hat{\kappa} = H + \frac{5}{3}b_4^2$. Therefore $J = \frac{5}{3}b_4^2$ and b_4 is a globally defined function on M^3 .

We want to show that the basis, we have constructed at each point p , can be extended differentiably to a neighborhood of the point p such that, at every point, K with respect to the frame $\{E, V, F\}$ has the previously described form.

Lemma 27. *Let M^3 be an affine hypersphere in $\mathbb{R}^4$ which admits a point-wise $\text{SO}(1, 1)$ -symmetry. Let $p \in M$. Then there exists a lightvector-frame $\{E, V, F\}$ defined in a neighborhood of the point p and a positive function b_4 such that K is given by:*

$$\begin{aligned}K(V, V) &= -2b_4V, & K(V, E) &= b_4E, & K(V, F) &= b_4F, \\ K(E, E) &= 0, & K(E, F) &= b_4V, & K(F, F) &= 0.\end{aligned}$$

Proof. First we want to show that at every point the vector $\mathbf{v}$ is uniquely defined and differentiable. We introduce a symmetric operator $\hat{A}$ by:

$$\widehat{\text{Ric}}(Y, Z) = h(\hat{A}Y, Z).$$

Clearly $\hat{A}$ is a differentiable operator on M^3 with $\hat{A}(\mathbf{e}) = 2(H + b_4^2)\mathbf{e}$, $\hat{A}(\mathbf{v}) = 2(H + 3b_4^2)\mathbf{v}$ and $\hat{A}(\mathbf{f}) = 2(H + b_4^2)\mathbf{f}$. Since $2(H + 3b_4^2) \neq 2(H + b_4^2)$,

the operator has two distinct eigenvalues. A standard result then implies that the eigendistributions are differentiable. We take V a local unit vectorfield spanning the 1-dimensional spacelike eigendistribution, and local orthonormal vectorfields $\tilde{T}$ and $\tilde{W}$ spanning the second eigendistribution. We define $E = \frac{1}{\sqrt{2}}(-\tilde{T} + \tilde{W})$ and $F = \frac{1}{\sqrt{2}}(\tilde{T} + \tilde{W})$. $\square$

Remark. It actually follows from the proof of the previous lemma that the vector field V is globally defined on M^3 .

In this section we always will work with the local frame constructed in the previous lemma. We denote the coefficients of the Levi-Civita connection with respect to this frame by:

$$\begin{aligned}\hat{\nabla}_E E &= a_{11}E + b_{11}V, & \hat{\nabla}_E V &= a_{12}E - b_{11}F, & \hat{\nabla}_E F &= -a_{12}V - a_{11}F, \\ \hat{\nabla}_V E &= a_{21}E + b_{21}V, & \hat{\nabla}_V V &= a_{22}E - b_{21}F, & \hat{\nabla}_V F &= -a_{22}V - a_{21}F, \\ \hat{\nabla}_F E &= a_{31}E + b_{31}V, & \hat{\nabla}_F V &= a_{32}E - b_{31}F, & \hat{\nabla}_F F &= -a_{32}V - a_{31}F.\end{aligned}$$

We will evaluate first the Codazzi and then the Gauss equations ((12) and (13)) to obtain more informations.

Lemma 28. *Let M^3 be an affine hypersphere in $\mathbb{R}^4$ which admits a pointwise $\text{SO}(1,1)$ -symmetry and $\{E, V, F\}$ the corresponding LVB. Then $0 = b_{11} = b_{21} = a_{22} = a_{32}$, $b_{31} = -a_{12}$ and $V(b_4) = -4a_{12}b_4$, $0 = E(b_4) = F(b_4)$.*

Proof. An evaluation of the Codazzi equations (12) leads to the following equations (they relate to eq1–eq5 and eq7–eq8 in the Mathematica notebook):

$$E(b_4) = -2b_{21}b_4, \quad 0 = b_{11}b_4, \quad (147)$$

$$0 = (a_{12} + b_{31})b_4, \quad 0 = E(b_4), \quad (148)$$

$$V(b_4) = -4a_{12}b_4, \quad (149)$$

$$0 = a_{32}b_4, \quad F(b_4) = 2a_{22}b_4 \quad V(b_4) = 4b_{31}b_4, \quad (150)$$

$$0 = F(b_4), \quad (151)$$

$$0 = a_{22}b_4, \quad (152)$$

$$0 = b_{21}b_4, \quad (153)$$

this proves the theorem. $\square$

An evaluation of the Gauss equations (13) leads to the following :

Lemma 29. *Let M^3 be an affine hypersphere in $\mathbb{R}^4$ which admits a pointwise $\text{SO}(1,1)$ -symmetry and $\{E, V, F\}$ the corresponding LVB. Then*

$$V(a_{12}) = -a_{12}^2 - 3b_4^2 - H, \quad (154)$$

$$F(a_{12}) = 0, \quad (155)$$

$$E(a_{12}) = 0, \quad (156)$$

$$E(a_{21}) - V(a_{11}) = a_{11}(a_{12} - a_{21}), \quad (157)$$

$$E(a_{31}) - F(a_{11}) = a_{12}^2 - 2a_{11}a_{31} - b_4^2 + H, \quad (158)$$

$$V(a_{31}) - F(a_{21}) = -a_{31}(a_{12} + a_{21}). \quad (159)$$

Proof. The equations relate to eq11–eq12 and eq14–eq16 in the Mathematica notebook. $\square$

As the vector field V is globally defined, we can define the distributions $L_1 = \text{span}\{V\}$ and $L_2 = \text{span}\{E, F\}$. In the following we will investigate these distributions.

Lemma 30. *The distribution L_1 is autoparallel (totally geodesic) with respect to $\hat{\nabla}$.*

Proof. From $\hat{\nabla}_V V = a_{22}E - b_{21}F = 0$ (cp. Lemma 28) the claim follows immediately. $\square$

Lemma 31. *The distribution L_2 is spherical with mean curvature normal $U_2 = -a_{12}V$.*

Proof. For $U_2 = -a_{12}V \in L_1 = L_2^\perp$ we have $h(\hat{\nabla}_{E_a} E_b, V) = h(E_a, E_b)h(U_2, V)$ for $E_a, E_b \in \{E, F\}$, and $h(\hat{\nabla}_{E_a} U_2, V) = h(-E_a(a_{12})V + a_{12}\hat{\nabla}_{E_a} V, V) = 0$ (cp. Lemma 28 and (156), (157)). $\square$

Remark. a_{12} depends of V but not of the choice of basis $\{E, F\}$ of L_2 . It therefore is a globally defined function on M^3 .

We introduce a coordinate function v by $\frac{\partial}{\partial v} := V$. Using the previous lemma, according to [PR93], we get:

Lemma 32. *(M^3, h) admits a warped product structure $M^3 = I \times_{ef} N^2$ with $f : I \rightarrow \mathbb{R}$ satisfying*

$$\frac{\partial f}{\partial v} = a_{12}. \quad (160)$$

Proof. Prop. 3 in [PR93] gives the warped product structure with warping function $\lambda_2 : I \rightarrow \mathbb{R}$. If we introduce $f = \ln \lambda_2$, following the proof we see that $-a_{12}V = U_2 = -\text{grad}(\ln \lambda_2) = -\text{grad } f$. $\square$

Lemma 33. *The curvature of N^2 is ${}^N K(N^2) = e^{2f}(-b_4^2 + a_{12}^2 + H)$.*

Proof. For the formulas cp. Lem. 16. Using the Gauss equation (13) (cp. the Mathematica notebook) we get:

$$(H - b_4^2)E = \hat{R}(E, F)E = {}^N\hat{R}(E, F)E - (E(a_{12})V + a_{12}(a_{12}E)) + h(E(a_{12})V + a_{12}(a_{12}E), E)F - h(F(a_{12})V + a_{12}(a_{12}F), E)E + a_{12}^2E.$$

$$\text{Thus } {}^N K(N^2) = \frac{{}^N h(-{}^N \hat{R}(E, F)E, F)}{{}^N h(E, E){}^N h(F, F) - {}^N h(E, F)^2} = e^{2f} h({}^N \hat{R}(E, F)E, F) = e^{2f}(-b_4^2 + a_{12}^2 + H). \quad \square$$

Summarized we have obtained the following structure equations (cp. (1), (2) and (3)):

$$D_V V = -2b_4 V + \xi, \quad (161)$$

$$D_V E = +(a_{21} + b_4)E, \quad (162)$$

$$D_V F = +(-a_{21} + b_4)F, \quad (163)$$

$$D_E V = +(a_{12} + b_4)E, \quad (164)$$

$$D_F V = +(a_{12} + b_4)F, \quad (165)$$

$$D_E E = +a_{11}E, \quad (166)$$

$$D_E F = -a_{11}F + (-a_{12} + b_4)V + \xi, \quad (167)$$

$$D_F E = +a_{31}E + (-a_{12} + b_4)V + \xi, \quad (168)$$

$$D_F F = -a_{31}F, \quad (169)$$

$$D_X \xi = -HX, \quad (170)$$

The Codazzi and Gauss equations ((12) and (13)) have the form (cp. Lem. 28 and 29):

$$V(b_4) = -4a_{12}b_4, \quad 0 = E(b_4) = F(b_4), \quad (171)$$

$$V(a_{12}) = -a_{12}^2 - 3b_4^2 - H, \quad F(a_{12}) = 0, \quad E(a_{12}) = 0, \quad (172)$$

$$E(a_{21}) - V(a_{11}) = a_{11}(a_{12} - a_{21}), \quad (173)$$

$$V(a_{31}) - F(a_{21}) = -a_{31}(a_{12} + a_{21}), \quad (174)$$

$$E(a_{31}) - F(a_{11}) = a_{12}^2 - 2a_{11}a_{31} - b_4^2 + H. \quad (175)$$

Our first goal is to find out how N^2 is immersed in $\mathbb{R}^4$, i. e. to find an immersion independent of v . A look at the structure equations (161) - (170) suggests to start with a linear combination of V and ξ .

We will solve the problem in two steps. First we look for a vector field X with $D_V X = \alpha X$ for some function α : We define $X := AV + \xi$ for some function A on M^3 . Then $D_V X = \alpha X$ iff $\alpha = A$ and $\frac{\partial}{\partial v} A = A^2 + 2b_4 A + H$, and $A := -a_{12} + b_4$ solves the latter differential equation. Next we want

to multiply X with some function β such that $D_V(\beta X) = 0$: We define a positive function β on $\mathbb{R}$ as the solution of the differential equation:

$$\frac{\partial}{\partial v}\beta = (a_{12} - b_4)\beta \quad (176)$$

with initial condition $\beta(v_0) > 0$. Then $D_V(\beta X) = 0$ and by (164), (170) and (165) we get (since β , a_{12} and b_4 only depend on v):

$$D_V(\beta((b_4 - a_{12})V + \xi)) = 0, \quad (177)$$

$$D_E(\beta((b_4 - a_{12})V + \xi)) = \beta(b_4^2 - a_{12}^2 - H)E, \quad (178)$$

$$D_F(\beta((b_4 - a_{12})V + \xi)) = \beta(b_4^2 - a_{12}^2 - H)F. \quad (179)$$

To obtain an immersion we need that $\nu := b_4^2 - a_{12}^2 - H$ vanishes nowhere, but we only get:

Lemma 34. *The function $\nu = b_4^2 - a_{12}^2 - H$ is globally defined, $\frac{\partial}{\partial v}(e^{2f}\nu) = 0$ and ν vanishes identically or nowhere on $\mathbb{R}$.*

Proof. Since $0 = \frac{\partial}{\partial v}^N K(N^2) = \frac{\partial}{\partial v}(e^{2f}(-\nu))$ (Lem. 33), we get by (160) $\frac{\partial}{\partial v}\nu = -2a_{12}\nu$. $\square$

8.1 The first case: $\nu \neq 0$ on M^3

We may, by translating f , i.e. by replacing N^2 with a homothetic copy of itself, assume that $e^{2f}\nu = \varepsilon_1$, where $\varepsilon_1 = \pm 1$.

Lemma 35. *$\Phi := \beta((b_4 - a_{12})V + \xi): M^3 \rightarrow \mathbb{R}^4$ induces the structure of an indefinite proper affine quadric, say $\tilde{\phi}$, mapping N^2 into a 3-dimensional linear subspace of $\mathbb{R}^4$.*

Proof. By (178) and (179) we have $\Phi_*(E_a) = \beta\nu E_a$ for $E_a \in \{E, F\}$. A further differentiation, using (166) (β and ν only depend on v), gives:

$$D_E\Phi_*(E) = \beta\nu D_E E = \beta\nu a_{11}E = a_{11}\Phi_*(E).$$

Similarly, we obtain the other derivatives, using (167) - (169), thus:

$$D_E\Phi_*(E) = a_{11}\Phi_*(E), \quad (180)$$

$$D_E\Phi_*(F) = -a_{11}\Phi_*(F) + e^{-2f}\varepsilon_1\Phi, \quad (181)$$

$$D_F\Phi_*(E) = a_{31}\Phi_*(E) + e^{-2f}\varepsilon_1\Phi, \quad (182)$$

$$D_F\Phi_*(F) = -a_{31}\Phi_*(F) \quad (183)$$

$$D_{E_a}\Phi = \beta e^{-2f}\varepsilon_1 E_a. \quad (184)$$

The foliation at $f = f_0$ gives an immersion of N^2 to M^3 , say π_{f_0} . Therefore, we can define an immersion of N^2 to $\mathbb{R}^4$ by $\tilde{\phi} := \Phi \circ \pi_{f_0}$, whose structure equations are exactly the equations above when $f = f_0$. Hence, we know

that $\tilde{\phi}$ maps N^2 into $\text{span}\{\Phi_*(E), \Phi_*(F), \Phi\}$, an affine hyperplane of $\mathbb{R}^4$ and $\frac{\partial}{\partial v}\Phi = 0$ (cp. (177)) implies $\Phi(v, x, y) = \tilde{\phi}(x, y)$. We can read off the coefficients of the difference tensor $K^{\tilde{\phi}}$ of $\tilde{\phi}$ (cf. (1) and (3)) and see that it vanishes. The affine metric introduced by this immersion corresponds with the metric on N^2 . Thus $\varepsilon_1\tilde{\phi}$ is the affine normal of $\tilde{\phi}$ and $\tilde{\phi}$ is an indefinite proper affine quadric with mean curvature ε_1 . $\square$

Our next goal is to find another linear combination of V and ξ , this time only depending on v . (Then we can express V in terms of ϕ and some function of v .)

Lemma 36. *Define $\delta := HV + (b_4 + a_{12})\xi$. Then there exist a constant vector $C \in \mathbb{R}^4$ and a function $a(v)$ such that*

$$\delta(v) = a(v)C.$$

Proof. Using (164) resp. (165) and (170) we obtain that $D_E\delta = 0 = D_F\delta$. Hence δ depends only on the variable v . Moreover, we get by (161), (172), (171) and (170) that

$$\begin{aligned} \frac{\partial}{\partial v}\delta &= D_V(HV + (b_4 + a_{12})\xi) \\ &= H(-2b_4V + \xi) + (-4a_{12}b_4 - a_{12}^2 - 3b_4^2 - H)\xi - (b_4 + a_{12})HV \\ &= (-3b_4 - a_{12})(HV + (b_4 + a_{12})\xi) \\ &= -(3b_4 + a_{12})\delta. \end{aligned}$$

This implies that there exists a constant vector C in $\mathbb{R}^4$ and a function $a(v)$ such that $\delta(v) = a(v)C$. $\square$

Notice that for an improper affine hypersphere ($H = 0$) ξ is constant and parallel to C . Combining $\tilde{\phi}$ and δ we obtain for V (cp. Lem. 35 and 36) that

$$V(v, x, y) = -\frac{a}{\nu}C + \frac{1}{\beta\nu}(b_4 + a_{12})\tilde{\phi}(x, y). \quad (185)$$

In the following we will use for the partial derivatives the abbreviation $\varphi_u := \frac{\partial}{\partial u}\varphi$, $u = v, x, y$.

Lemma 37.

$$\begin{aligned} \varphi_v &= -\frac{a}{\nu}C + \frac{\partial}{\partial v}\left(\frac{1}{\beta\nu}\right)\tilde{\phi}, \\ \varphi_x &= \frac{1}{\beta\nu}\tilde{\phi}_x, \\ \varphi_y &= \frac{1}{\beta\nu}\tilde{\phi}_y. \end{aligned}$$

Proof. As by (176) and Lem. 34 $\frac{\partial}{\partial v} \frac{1}{\beta\nu} = \frac{1}{\beta\nu}(b_4 + a_{12})$, we obtain the equation for $\varphi_v = V$ by (185). The other equations follow from (178) and (179). $\square$

It follows by the uniqueness theorem of first order differential equations and applying a translation that we can write

$$\varphi(v, x, y) = \tilde{a}(v)C + \frac{1}{\beta\nu}(v)\tilde{\phi}(x, y)$$

for a suitable function $\tilde{a}$ depending only on the variable v . Since C is transversal to the image of $\tilde{\phi}$ (cp. Lem. 35 and 36, $\nu \neq 0$), we obtain that after applying an equiaffine transformation we can write: $\varphi(v, x, y) = (\gamma_1(v), \gamma_2(v)\phi(x, y))$, where $\tilde{\phi}(x, y) = (0, \phi(x, y))$. Thus we have proven the following:

Theorem 17. *Let M^3 be an indefinite affine hypersphere of $\mathbb{R}^4$ which admits a pointwise $SO(1, 1)$ -symmetry. Let $b_4^2 - a_{12}^2 \neq H$ for some $p \in M^3$. Then M^3 is affine equivalent to*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (v, x, y) \mapsto (\gamma_1(v), \gamma_2(v)\phi(x, y)),$$

where $\phi : N^2 \rightarrow \mathbb{R}^3$ is a one-sheeted hyperboloid and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve.

Since the immersion φ has the same form as in the case of $SO(2)$ -symmetry, most of the computations for the converse theorem are the same (cp. Theorem 5 and the computations thereafter). We just have to keep track of the different coordinates and that now the affine metric g of ϕ is indefinite. If $H \neq 0$ then φ is indefinite and V is spacelike iff $H \text{sign}(\gamma_1\gamma_2' - \gamma_1'\gamma_2) = \text{sign}(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2')$. If $H = 0$, then φ is indefinite and V is spacelike iff $\text{sign}(\gamma_2') = \text{sign}(\gamma_1'\gamma_2'' - \gamma_1''\gamma_2')$. Also the computation of K is completely the same as before in the proof of Theorem 6, implying that K_{φ_v} restricted to the space spanned by φ_x and φ_y is a multiple of the identity. Taking V in direction of φ_v , we can construct a LVB $\{E, V, F\}$ with E, F spanning $\text{span}\{\varphi_x, \varphi_y\}$ such that $b_4 \neq 0$ and $b_1 = b_2 = b_5 = b_6 = 0$. By the considerations in Sec. 4 we see that φ admits a pointwise $SO(1, 1)$ -symmetry.

Theorem 18. *Let $\phi : N^2 \rightarrow \mathbb{R}^3$ be a one-sheeted hyperboloid and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve, such that $\varphi(t, v, w) = (\gamma_1(t), \gamma_2(t)\phi(v, w))$ defines a 3-dimensional indefinite affine hypersphere. Then $\varphi(N^2 \times I)$ admits a pointwise $SO(1, 1)$ -symmetry.*

- (i) *If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_2^2|\gamma_1\gamma_2' - \gamma_1'\gamma_2|^5 = |\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'|(\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.*
- (ii) *If $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_2^2|\gamma_2'|^5 = |\gamma_1'\gamma_2'' - \gamma_1''\gamma_2'|(\gamma_1')^2 \neq 0$, then φ defines a 3-dimensional indefinite improper affine hypersphere.*

8.1.1 The second case: $\nu \equiv 0$ and $H \neq 0$ on M^3

Next, we consider the case that $H = b_4^2 - a_{12}^2$ and $H \neq 0$ on M^3 . It follows that $b_4 \neq \pm a_{12}$ on M^3 .

We already have seen that M^3 admits a warped product structure. For a fixed point v_0 , we get from (166) - (169), (178) and (179), using the notation $\tilde{\xi} = (b_4 - a_{12})V + \xi$:

$$D_E E = + a_{11} E, \quad (186)$$

$$D_E F = - a_{11} F + \tilde{\xi}, \quad (187)$$

$$D_F E = + a_{31} E + \tilde{\xi}, \quad (188)$$

$$D_F F = - a_{31} F, \quad (189)$$

$$D_{E_a} \tilde{\xi} = 0, \quad E_a \in \{E, F\}. \quad (190)$$

Thus, if x and y are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(v_0, x, y)$ as a indefinite improper affine quadric, i. e. a hyperbolic paraboloid, in a 3-dimensional linear subspace.

After applying a translation and a change of coordinates, we may assume that

$$\varphi(v_0, x, y) = (x, y, f(x, y), 0), \quad (191)$$

with affine normal $\tilde{\xi}(v_0, x, y) = (0, 0, 1, 0)$. Notice that by (177) - (179) we have that $D_X(\beta\tilde{\xi}) = 0$, $X \in \mathcal{X}(M)$. Taking suitable initial conditions for the function β ($\beta(v_0) = 1$), we get that $\beta\tilde{\xi} = (0, 0, 1, 0)$ and thus

$$\xi = \frac{1}{\beta}(0, 0, 1, 0) - (b_4 - a_{12})V. \quad (192)$$

Applied to (161) we get for fixed x, y the following second order linear ODE:

$$D_V V = (a_{12} - 3b_4)V + \frac{1}{\beta}(0, 0, 1, 0).$$

The solution has the form

$$V(v, x, y) = \delta_1(v)(\delta_2(v)(0, 0, 1, 0) + C(x, y)), \quad (193)$$

where $\delta'_1(v) = (a_{12} - 3b_4)\delta_1(v)$ and $\delta'_2(v) = \frac{1}{\beta\delta_1}(v)$. To obtain V at v_0 , we consider (164) and (165) and get that

$$D_{E_a}(V - (a_{12} + b_4)\varphi) = 0, \quad E_a \in \{E, F\}.$$

Evaluating at $v = v_0$, this means that there exists a constant vector $\tilde{C}$ such that

$$V(v_0, x, y) = (a_{12} + b_4)(v_0)\varphi(v_0, x, y) + \tilde{C}. \quad (194)$$

Since $\tilde{C}$ must be transversal to $\text{span}\{E, F, \varphi\} = \text{span}\{E, F, \xi\}$, we can write:

$$V(v_0, x, y) = \alpha_1(x, y, f(x, y), \alpha_2),$$

where $\alpha_1, \alpha_2 \neq 0$ and we applied an equiaffine transformation so that $\tilde{C} = (0, 0, 0, \alpha_1 \alpha_2)$. If we compare with (193), then we see that $C(x, y) = \frac{\alpha_1}{\delta_1(v_0)}(x, y, f(x, y), \alpha_2) - \delta_2(v_0)(0, 0, 1, 0)$, and therefore:

$$V(v, x, y) = \delta_1(v)(x, y, f(x, y) + \delta_2(v), \alpha_2), \quad (195)$$

where $\delta'_1(v) = (a_{12} - 3b_4)\delta_1(v)$, $\delta_1(v_0) = \alpha_1$, and $\delta'_2(v) = \frac{1}{\beta\delta_1}(v)$, $\delta_2(v_0) = 0$. Respecting the initial condition (191), after integration we obtain that

$$\varphi(v, x, y) = (\gamma_1(v)x, \gamma_1(v)y, \gamma_1(v)f(x, y) + \gamma_2(v), \alpha_2(\gamma_1(v) - 1)),$$

where $\gamma'_1(v) = \delta_1(v)$, $\gamma_1(v_0) = 1$, and $\gamma'_2(v) = \delta_1(v)\delta_2(v)$, $\gamma_2(v_0) = 0$. After applying an affine transformation we have shown:

Theorem 19. *Let M^3 be an indefinite proper affine hypersphere of $\mathbb{R}^4$ which admits a pointwise $SO(1, 1)$ -symmetry. Let $H = b_4^2 - a_{12}^2 (\neq 0)$ on M^3 . Then M^3 is affine equivalent with*

$$\varphi : I \times N^2 \rightarrow \mathbb{R}^4 : (v, x, y) \mapsto (\gamma_1(v)x, \gamma_1(v)y, \gamma_1(v)f(x, y) + \gamma_2(v), \gamma_1(v)),$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (x, y) \mapsto (x, y, f(x, y))$ is a hyperbolic paraboloid with affine normal $(0, 0, 1)$ and $\gamma : I \rightarrow \mathbb{R}^2$ is a curve.

Since the immersion φ has the same form as in the case of $SO(2)$ -symmetry, most of the computations for the converse theorem are the same (cp. Theorem 7 and the computations thereafter). We just have to keep track of the different coordinates and that ψ is indefinite, i. e. $f_{xx}f_{yy} - f_{xy}^2 = -1$. Thus φ is indefinite and V is spacelike iff $H \text{sign}(\gamma_1\gamma'_2 - \gamma'_1\gamma_2) = \text{sign}(\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2)$. Also the computation of K is completely the same as before in the proof of Theorem 8, implying that K_{φ_v} restricted to the space spanned by φ_x and φ_y is a multiple of the identity. Taking V in direction of φ_v , we can construct a LVB $\{E, V, F\}$ with E, F spanning $\text{span}\{\varphi_x, \varphi_y\}$ such that $b_4 \neq 0$ and $b_1 = b_2 = b_5 = b_6 = 0$. By the considerations in Sec. 4 we see that φ admits a pointwise $SO(1, 1)$ -symmetry.

Theorem 20. *Let $\psi : N^2 \rightarrow \mathbb{R}^3$ be a hyperbolic paraboloid with affine normal $(0, 0, 1)$, and let $\gamma : I \rightarrow \mathbb{R}^2$ be a curve, such that $\varphi(v, x, y) = (\gamma_1(v)x, \gamma_1(v)y, \gamma_1(v)f(x, y) + \gamma_2(v), \gamma_1(v))$ defines a 3-dimensional indefinite proper affine hypersphere. Then $\varphi(N^2 \times I)$ admits a pointwise $SO(1, 1)$ -symmetry.*

Moreover, if $\gamma = (\gamma_1, \gamma_2)$ satisfies $\gamma_1^2|\gamma_1\gamma'_2 - \gamma'_1\gamma_2|^5 = |\gamma'_1\gamma''_2 - \gamma''_1\gamma'_2|(\gamma'_1)^2 \neq 0$, then φ defines a 3-dimensional indefinite proper affine hypersphere.

8.1.2 The third case: $\nu \equiv 0$ and $H = 0$ on M^3

The final cases now are that $\nu \equiv 0$ and $H = 0$ on the whole of M^3 and hence $a_{12} = \pm b_4$.

First we consider the case that $a_{12} = b_4 =: b > 0$. Again we use that M^3 admits a warped product structure and we fix a parameter v_0 . At the point v_0 , we have by (166)–(170):

$$\begin{aligned} D_E E &= +a_{11} E, \\ D_E F &= -a_{11} F + \xi, \\ D_F E &= +a_{31} E + \xi, \\ D_F F &= -a_{31} F, \\ D_X \xi &= 0. \end{aligned}$$

Thus, if x and y are local coordinates which span the second distribution L_2 , then we can interpret $\varphi(v_0, x, y)$ as an indefinite improper affine quadric, i. e. a hyperbolic paraboloid, in a 3-dimensional linear subspace.

Completely analogous to the case of $SO(2)$ -symmetry we obtain by (171) that we can take $b = \frac{1}{4v}$, $v > 0$, and arrive at the following linear second order ordinary differential equation (cp. (86)):

$$\frac{\partial^2}{\partial v^2} \varphi + \frac{1}{2v} \frac{\partial}{\partial v} \varphi = -\xi,$$

where $\xi(v, x, y) = (0, 0, 1, 0)$. Thus we end up with (cp. (87)):

$$\varphi(v, x, y) = (vx, vy, vf(x, y) - cv^4, v), \quad c, v > 0. \quad (196)$$

Next we consider the case that $-a_{12} = b_4 =: b > 0$. Again we use that M^3 admits a warped product structure and we fix a parameter v_0 . As in the second case we can interpret $\varphi(v_0, x, y)$ as an indefinite improper affine quadric (cp. (186)–(189)) with constant affine normal $\tilde{\xi} = 2bV + \xi$, i. e. we get a hyperbolic paraboloid in a 3-dimensional linear subspace. Also, in analogy to the case of $SO(2)$ -symmetry (just notice the different sign in the definition of $\tilde{\xi}$), we can take $b = \frac{1}{4|v|}$, $v < 0$, and arrive at the following ordinary differential equation:

$$\frac{\partial}{\partial v} \varphi = \sqrt{|v|} C + 2v\xi, \quad v < 0.$$

As before we get that the hypersurface is affine equivalent to:

$$\varphi(v, x, y) = (x, y, f(x, y) + cv^3, v^4), \quad c, v > 0. \quad (197)$$

Combining both results (196) and (197) we have:

Theorem 21. *Let M^3 be an indefinite improper affine hypersphere of $\mathbb{R}^4$ which admits a pointwise $SO(1, 1)$ -symmetry. Let $a_{12}^2 = b_4^2$ on M^3 . Then M^3 is affine equivalent with either*

$$\begin{aligned} \varphi : I \times N^2 &\rightarrow \mathbb{R}^4 : (v, x, y) \mapsto (vx, vy, vf(x, y) - cv^4, v), \quad (a_{12} = b_4) \quad \text{or} \\ \varphi : I \times N^2 &\rightarrow \mathbb{R}^4 : (v, x, y) \mapsto (x, y, f(x, y) + cv^3, v^4), \quad (-a_{12} = b_4) \end{aligned}$$

where $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ is a hyperbolic paraboloid with affine normal $(0, 0, 1)$ and $c, t \in \mathbb{R}^+$.

The computations for the converse statement can be done completely analogous to the previous cases, they even are simpler (the curve is given parametrized).

Theorem 22. *Let $\psi : N^2 \rightarrow \mathbb{R}^3 : (v, w) \mapsto (v, w, f(v, w))$ be a hyperbolic paraboloid with affine normal $(0, 0, 1)$. Define $\varphi(t, v, w) = (tv, tw, tf(v, w) - ct^4, t)$ or $\varphi(t, v, w) = (v, w, f(v, w) + ct^3, t^4)$, where $t \in \mathbb{R}^+$, $c \neq 0$. Then φ defines a 3-dimensional indefinite improper affine hypersphere, which admits a pointwise $SO(1, 1)$ -symmetry.*

References

- [Bry01] R. L. Bryant, *Second order families of special lagrangian 3-folds*, preprint, arXiv:math.DG/0007128 v2, 2001.
- [Che93] B.-Y. Chen, *Some pinching and classification theorems for minimal submanifolds*, Arch. Math. (Basel) **60** (1993), 568–578.
- [Dil89] F. Dillen, *Equivalence theorems in affine differential geometry*, Geom. Dedicata **32** (1989), 81–92.
- [DMV00] F. Dillen, M. Magid, and L. Vrancken, *Affine hyperspheres with constant affine sectional curvature*, Geometry and topology of submanifolds, X (Beijing/Berlin, 1999) (River Edge, NJ), World Sci. Publishing, River Edge, NJ, 2000, pp. 31–53.
- [DNV90] F. Dillen, K. Nomizu, and L. Vrancken, *Conjugate connections and Radon’s theorem in affine differential geometry*, Monatsh. Math. **109** (1990), 221–235.
- [Gre67] W. H. Greub, *Linear algebra*, Third edition. Die Grundlehren der Mathematischen Wissenschaften, Band 97, Springer-Verlag New York, Inc., New York, 1967.
- [JL05] F. Jia and A.-M. Li, *Locally strongly convex hypersurfaces with constant affine mean curvature*, Differential Geom. Appl. **22** (2005), 199–214.
- [KSV01] M. Kriele, C. Scharlach, and L. Vrancken, *An extremal class of 3-dimensional elliptic affine spheres*, Hokkaido Math. J. **30** (2001), 1–23.

- [KV99] M. Kriele and L. Vrancken, *An extremal class of three-dimensional hyperbolic affine spheres*, Geom. Dedicata **77** (1999), 239–252, MR 2000h:53016, ZBL 991.00978.
- [LSZ93] A. M. Li, U. Simon, and G. S. Zhao, *Global affine differential geometry of hypersurfaces*, de Gruyter Expositions in Mathematics, vol. 11, Walter de Gruyter & Co., Berlin, 1993.
- [LS] Y. Lu and C. Scharlach, *Affine hypersurfaces admitting a point-wise symmetry*, submitted, arXiv:math.DG/0510150.
- [MR92] M. A. Magid and P. J. Ryan, *Affine 3-spheres with constant affine curvature*, Trans. Am. Math. Soc. **330** (1992), 887–901.
- [Nöl96] S. Nölker, *Isometric immersions of warped products*, Differential Geom. Appl. **6** (1996), 1–30.
- [NS94] K. Nomizu and T. Sasaki, *Affine differential geometry*, Cambridge Tracts in Mathematics, vol. 111, Cambridge University Press, Cambridge, 1994, Geometry of affine immersions.
- [O’N83] B. O’Neill, *Semi-Riemannian geometry*, Pure and Applied Mathematics, vol. 103, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1983, With applications to relativity.
- [PR93] R. Ponge and H. Reckziegel, *Twisted products in pseudo-Riemannian geometry*, Geom. Dedicata **48** (1993), 15–25.
- [SSVV97] C. Scharlach, U. Simon, L. Verstraelen, and L. Vrancken, *A new intrinsic curvature invariant for centroaffine hypersurfaces*, Beitrüge Algebra Geom. **38** (1997), 437–458.
- [SV96] C. Scharlach and L. Vrancken, *A curvature invariant for centroaffine hypersurfaces. II*, Geometry and topology of submanifolds, VIII (Brussels, 1995/Nordfjordeid, 1995) (River Edge, NJ), World Sci. Publishing, River Edge, NJ, 1996, pp. 341–350.
- [Vra00] L. Vrancken, *The Magid-Ryan conjecture for equiaffine hyperspheres with constant sectional curvature*, J. Differential Geom. **54** (2000), 99–138.
- [Vra04] ———, *Special classes of three dimensional affine hyperspheres characterized by properties of their cubic form*, Contemporary geometry and related topics (River Edge, NJ), World Sci. Publishing, River Edge, NJ, 2004, pp. 431–459.

Authors address: C. Scharlach, Technische Universität Berlin, Fakultät II,
Inst. f. Mathematik, MA 8-3, 10623 Berlin, Germany
E-mail: schar@math.tu-berlin.de